

PARABOLIC AND ELLIPTIC SYSTEMS WITH VMO COEFFICIENTS

HONGJIE DONG AND DOYOON KIM

ABSTRACT. We consider second order parabolic and elliptic systems with leading coefficients having the property of vanishing mean oscillation (VMO) in the spatial variables. An $L_q - L_p$ theory is established for systems both in divergence and non-divergence form. Higher order parabolic and elliptic systems are also discussed briefly.

1. INTRODUCTION

The L_p theory of second order parabolic and elliptic equations has been studied extensively by many authors for more than fifty years. It is of particular interest not only because of its various important applications in nonlinear equations, but also due to its subtle links with the theory of stochastic processes. For scalar equations, the solvability theory in L_p spaces has been well established; see, for example, [7], [8], [2], [15]-[20], [3] and references therein.

In contrary, until quite recently, there are very few results of L_p theory for parabolic systems with discontinuous coefficients. On the other hand, quite naturally, many evolutionary equations arising from physical and economical problems are coupled systems instead of scalar equations, such as the Navier-Stokes equations.

In this paper we consider two types of parabolic operators

$$P\mathbf{u} = -\mathbf{u}_t + A^{\alpha\beta}D_{\alpha\beta}\mathbf{u} + B^\alpha D_\alpha\mathbf{u} + C\mathbf{u}, \quad (1)$$

$$\mathcal{P}\mathbf{u} = -\mathbf{u}_t + D_\alpha(A^{\alpha\beta}D_\beta\mathbf{u} + \hat{B}^\alpha\mathbf{u}) + B^\alpha D_\alpha\mathbf{u} + C\mathbf{u} \quad (2)$$

acting on (column) vector-valued functions $\mathbf{u} = (u^1, \dots, u^m)^T$ given on

$$\mathbb{R}^{d+1} = \{(t, x) : t \in \mathbb{R}, x = (x_1, \dots, x_d) \in \mathbb{R}^d\}.$$

Here, we have used the notations

$$D_\alpha\mathbf{u} = \mathbf{u}_{x_\alpha}, \quad D_{\alpha\beta}\mathbf{u} = \mathbf{u}_{x_\alpha x_\beta} \quad (\alpha, \beta = 1, \dots, d)$$

and the usual summation conventions over repeated indices are assumed. The coefficients $A^{\alpha\beta}$, B^α , \hat{B}^α , and C are $m \times m$ matrix-valued functions given on \mathbb{R}^{d+1} ; i.e., $A^{\alpha\beta} = [A_{ij}^{\alpha\beta}(t, x)]_{m \times m}$, etc. When the coefficients $A^{\alpha\beta}$, B^α , \hat{B}^α and C^α are independent of t , we also define and consider elliptic operators in divergence and non-divergence form

$$L\mathbf{u} = A^{\alpha\beta}D_{\alpha\beta}\mathbf{u} + B^\alpha D_\alpha\mathbf{u} + C\mathbf{u}, \quad (3)$$

$$\mathcal{L}\mathbf{u} = D_\alpha(A^{\alpha\beta}D_\beta\mathbf{u} + \hat{B}^\alpha\mathbf{u}) + B^\alpha D_\alpha\mathbf{u} + C\mathbf{u} \quad (4)$$

acting on vector valued functions $\mathbf{u} = (u^1, \dots, u^m)^T$ given on \mathbb{R}^d .

H. Dong was partially supported by a start-up funding from the Division of Applied Mathematics of Brown University and NSF grant number DMS 0800129.

We note that a similar problem is treated in a recent paper [14], where the authors considered higher order parabolic systems in non-divergence form. Under the assumption that leading coefficients are bounded, time-independent, and VMO in the spatial variables, the solvability is established in $L_q(L_p)$ spaces with A_p Muckenhoupt weights, by using estimates of integral operators of Calderon-Zygmund type and related commutators with BMO functions.

It turns out that, until recently, [14] is one of the few results on the L_p solvability of parabolic systems with discontinuous coefficients. With continuous leading coefficients, similar results can be found in an earlier paper [9]. For local L_p estimates of elliptic systems in non-divergence form with VMO coefficients, we refer interested readers to [25]. We also would like to bring attention to an interesting paper [4], in which the authors obtain $W^{1,p}$ solvability of a conormal derivative problem of elliptic systems in divergence form with VMO coefficients in domains satisfying a mild geometric condition. For other regularity results for elliptic and parabolic systems, see [26], [5], [1], [13], [10], [22], [23], [24], [6] and references therein.

In this paper we consider parabolic and elliptic operators both in divergence and non-divergence form with leading coefficients merely measurable in the time variable and VMO in the spatial variables. This is the same class of coefficients used in [19] and [20], and is denoted as VMO_x (see the definition in Section 3), which is strictly larger than those considered in [23] and [14]. Moreover, we assume that the leading coefficients satisfy the Legendre-Hadamard condition, which is weaker and more natural than the usual strong ellipticity condition (see Assumption 2.1).

Under these assumptions, we establish $L_q(L_p)$ solvability of both divergence and non-divergence form parabolic systems (cf. Theorem 2.3 and 2.4) extending the corresponding results for scalar equations in [20]. As a corollary, we obtain L_p solvability for divergence and non-divergence form elliptic systems as well (see Theorem 2.5 and 2.6).

Our approach is based on a method from [19] and [20]. Unlike the arguments in [7], [8], [14], [25] and [22]-[24], which are based on certain estimates of Calderon-Zygmund theorem and the Coifman-Rochberg-Weiss commutator theorem, our proofs rely mainly on pointwise estimates of sharp functions of second spatial derivatives of solutions to systems as in [19], [20]; see also [16] and [17].

Roughly speaking, there are three steps in the proofs of our main theorems. In the *first* step, we consider the systems with coefficients depending on the time variable only. By using the Fourier transform methods, we prove the solvability in L_2 space, from which we establish an estimate for sharp functions of derivatives of solutions to systems. Applying the Fefferman-Stein theorem on sharp functions and the Hardy-Littlewood maximal function theorem, we get the solvability in L_p spaces for $p > 2$, and then in L_p spaces for $p < 2$ via a duality argument. We also get another sharp function estimate from this solvability theorem. In the *second* step, we employ the sharp function estimate obtained in the previous step and a perturbation argument to obtain the L_p solvability of systems with leading coefficients in VMO_x . Finally in the *third* step, adapting an approach suggested in [18], we arrive at the $L_q(L_p)$ solvability for $q \geq p$ in the non-divergence form and without this restriction in the divergence form.

To the best of our knowledge, the question of the $L_q(L_p)$ solvability in the non-divergence form for $q < p$ with VMO_x coefficients is still open even for scalar equations. We remark that such restriction is not imposed in [14]. However, our

results are not covered by those in [14] because the coefficients are assumed to be time-independent in [14].

The paper is organized as follows. The main results, Theorem 2.3, 2.4, 2.5, and 2.6 are stated in the next section. First we consider systems in non-divergence form. Section 3 is devoted to some preliminary estimates including Theorem 3.1, which may be considered a preliminary version of Theorem 2.3. Theorem 3.1 is about the solvability of parabolic systems with coefficients independent of the spatial variables and is proved in Section 4. Next, Theorem 5.1 is proved in Section 5 as another special case of Theorem 2.3 when $p = q$ (see also Remark 5.2). The main tool of the proof is a pointwise estimate of the sharp function of second derivatives (Lemma 5.3). With these preparations, we are able to prove Theorem 2.3 in Section 6. In the remaining part of the paper, we turn to consider systems in divergence form. In Section 7, we state and prove Theorem 7.1, which is a special case of Theorem 2.4 when $p = q$, and is a counterpart of Theorem 5.1. Then Theorem 2.5 is proved in Section 8. Finally, the results of L_p theory of elliptic systems, i.e., Theorem 2.5 and 2.6, are derived in Section 9 by using Theorem 2.3 and 2.4. Higher order elliptic and parabolic systems are also discussed without many details at the end of this paper.

We finish the section by introducing some notation. Throughout the paper, we always assume that $1 < p, q < \infty$ unless explicitly specified otherwise. By $N(d, p, \dots)$ we mean that N is a constant depending only on the prescribed quantities d, p, \dots . For a (matrix-valued) function $F(t, x)$ in \mathbb{R}^{d+1} , we set

$$(F)_{\mathcal{D}} = \frac{1}{|\mathcal{D}|} \int_{\mathcal{D}} F(t, x) dx dt = \int_{\mathcal{D}} F(t, x) dx dt,$$

where \mathcal{D} is an open subset in \mathbb{R}^{d+1} and $|\mathcal{D}|$ is the $d + 1$ -dimensional Lebesgue measure of \mathcal{D} . For $-\infty \leq S < T \leq \infty$, we set

$$L_{q,p}((S, T) \times \mathbb{R}^d) = L_q((S, T), L_p(\mathbb{R}^d)),$$

i.e., $F(t, x) \in L_{q,p}((S, T) \times \mathbb{R}^d)$ if

$$\|F\|_{L_{q,p}((S,T) \times \mathbb{R}^d)} = \left(\int_S^T \left(\int_{\mathbb{R}^d} |F(t, x)|^p dx \right)^{q/p} dt \right)^{1/q} < \infty.$$

$$W_{q,p}^{1,2}((S, T) \times \mathbb{R}^d) = \{u : u, u_t, Du, D^2u \in L_{q,p}((S, T) \times \mathbb{R}^d)\},$$

$$W_p^{1,2}((S, T) \times \mathbb{R}^d) = W_{p,p}^{1,2}((S, T) \times \mathbb{R}^d),$$

$$\mathcal{H}_{q,p}^1((S, T) \times \mathbb{R}^d) = (1 - \Delta)^{1/2} W_{q,p}^{1,2}((S, T) \times \mathbb{R}^d),$$

$$\mathcal{H}_p^1((S, T) \times \mathbb{R}^d) = \mathcal{H}_{p,p}^1((S, T) \times \mathbb{R}^d),$$

$$\mathbb{H}_{q,p}^{-1}((S, T) \times \mathbb{R}^d) = (1 - \Delta)^{1/2} L_{q,p}((S, T) \times \mathbb{R}^d),$$

$$\mathbb{H}_p^{-1}((S, T) \times \mathbb{R}^d) = \mathbb{H}_{p,p}^{-1}((S, T) \times \mathbb{R}^d).$$

For any $T \in (-\infty, \infty]$, we denote

$$\mathbb{R}_T = (-\infty, T), \quad \mathbb{R}_T^{d+1} = \mathbb{R}_T \times \mathbb{R}^d.$$

2. MAIN RESULTS

Assumption 2.1. The matrices $A^{\alpha\beta}$, B^α , \hat{B}^α and C satisfy

$$|A^{\alpha\beta}(t, x)| \leq K, \quad |B^\alpha(t, x)| \leq K, \quad |\hat{B}^\alpha(t, x)| \leq K, \quad |C(t, x)| \leq K,$$

$$A_{ij}^{\alpha\beta}(t, x) \xi_\alpha \xi_\beta \vartheta^i \vartheta^j \geq \delta |\xi|^2 |\vartheta|^2$$

for all $(t, x) \in \mathbb{R}^{d+1}$, $\xi \in \mathbb{R}^d$, $\vartheta \in \mathbb{R}^m$, for some constants $0 < \delta < 1$ and $K > 0$.

Another assumption on $A = [A_{ij}^{\alpha\beta}]$ is that they are in the class of VMO_x , that is, $A_{ij}^{\alpha\beta}(t, x)$ are measurable in $t \in \mathbb{R}$ and VMO in $x \in \mathbb{R}^d$. A precise description of this assumption is given below using the following notations. Let

$$B_r(x) = \{y \in \mathbb{R}^d : |x - y| < r\}, \quad Q_r(t, x) = (t - r^2, t) \times B_r(x).$$

Set $B_r = B_r(0)$ and $|B_r|$ to be the d -dimensional volume of B_r . Also set

$$\begin{aligned} \text{osc}_x \left(A_{ij}^{\alpha\beta}, Q_r(t, x) \right) \\ = r^{-2} |B_r|^{-2} \int_{t-r^2}^t \int_{y, z \in B_r(x)} \left| A_{ij}^{\alpha\beta}(s, y) - A_{ij}^{\alpha\beta}(s, z) \right| dy dz ds \end{aligned}$$

and

$$A_R^\# = \sup_{(t, x) \in \mathbb{R}^{d+1}} \sup_{r \leq R} \sum_{\alpha, \beta=1}^d \sum_{i, j=1}^m \text{osc}_x \left(A_{ij}^{\alpha\beta}, Q_r(t, x) \right).$$

In case A is independent of t , naturally we set

$$\begin{aligned} \text{osc} \left(A_{ij}^{\alpha\beta}, B_r(x) \right) &= |B_r|^{-2} \int_{y, z \in B_r(x)} \left| A_{ij}^{\alpha\beta}(y) - A_{ij}^{\alpha\beta}(z) \right| dy dz, \\ A_R^\# &= \sup_{x \in \mathbb{R}^d} \sup_{r \leq R} \sum_{\alpha, \beta=1}^d \sum_{i, j=1}^m \text{osc} \left(A_{ij}^{\alpha\beta}, B_r(x) \right). \end{aligned}$$

Assumption 2.2. There is an increasing continuous function $\omega(r)$ defined on $[0, \infty)$ such that $\omega(0) = 0$ and $A_R^\# \leq \omega(R)$.

Now we state our main results. In Theorem 2.3 – 2.6, we use the notations $\mathbf{f} = (f^1, \dots, f^m)^T$, $\mathbf{g} = (\mathbf{g}_\alpha)$, $\mathbf{g}_\alpha = (g_\alpha^1, \dots, g_\alpha^m)^T$, $\alpha = 1, \dots, d$.

Theorem 2.3. *Let $1 < p \leq q < \infty$, $0 < T < \infty$, and the coefficient matrices of P satisfy Assumption 2.1 and 2.2. Then for any $\mathbf{f} \in L_{q,p}((0, T) \times \mathbb{R}^d)$, there exists a unique $\mathbf{u} \in W_{q,p}^{1,2}((0, T) \times \mathbb{R}^d)$ such that $P\mathbf{u} = \mathbf{f}$ in $(0, T) \times \mathbb{R}^d$ and $\mathbf{u}(0, \cdot) = 0$. Furthermore, there is a constant N , depending only on d, m, p, q, δ, K, T , and the function ω , such that*

$$\|\mathbf{u}\|_{W_{q,p}^{1,2}((0, T) \times \mathbb{R}^d)} \leq N \|P\mathbf{u}\|_{L_{q,p}((0, T) \times \mathbb{R}^d)}$$

for any $\mathbf{u} \in W_{q,p}^{1,2}((0, T) \times \mathbb{R}^d)$ satisfying $\mathbf{u}(0, \cdot) = 0$.

Theorem 2.4. *Let $p, q \in (1, \infty)$, $T \in (0, \infty)$, and the coefficient matrices of \mathcal{P} satisfy Assumption 2.1 and 2.2. Then for any $\mathbf{f}, \mathbf{g}_\alpha \in L_{q,p}((0, T) \times \mathbb{R}^d)$, there exists a unique $\mathbf{u} \in \mathcal{H}_{q,p}^1((0, T) \times \mathbb{R}^d)$ such that $\mathcal{P}\mathbf{u} = \mathbf{f} + D_\alpha \mathbf{g}_\alpha$ in $(0, T) \times \mathbb{R}^d$ and $\mathbf{u}(0, \cdot) = 0$. Furthermore, there is a constant N , depending only on d, m, p, q, δ, K, T , and the function ω , such that*

$$\|\mathbf{u}\|_{\mathcal{H}_{q,p}^1((0, T) \times \mathbb{R}^d)} \leq N (\|\mathbf{f}\|_{L_{q,p}((0, T) \times \mathbb{R}^d)} + \|\mathbf{g}\|_{L_{q,p}((0, T) \times \mathbb{R}^d)}).$$

Theorem 2.5. *Let $p \in (1, \infty)$, and the coefficient matrices of L satisfy Assumption 2.1, 2.2. Then there exist constants $\lambda_0 \geq 0$ and N , depending only on d, m, p, δ, K , and the function ω , such that*

$$\lambda \|\mathbf{u}\|_{L_p(\mathbb{R}^d)} + \sqrt{\lambda} \|D\mathbf{u}\|_{L_p(\mathbb{R}^d)} + \|D^2\mathbf{u}\|_{L_p(\mathbb{R}^d)} \leq N \|L\mathbf{u} - \lambda\mathbf{u}\|_{L_p(\mathbb{R}^d)} \quad (5)$$

holds for any $\mathbf{u} \in W_p^2(\mathbb{R}^d)$ and $\lambda \geq \lambda_0$. Moreover, for any $\lambda > \lambda_0$ and $\mathbf{f} \in L_p(\mathbb{R}^d)$, there exists a unique $\mathbf{u} \in W_p^2(\mathbb{R}^d)$ satisfying

$$L\mathbf{u} - \lambda\mathbf{u} = \mathbf{f} \quad \text{in } \mathbb{R}^d.$$

Theorem 2.6. *Let $p \in (1, \infty)$ and the coefficient matrices of \mathcal{L} satisfy Assumption 2.1 and 2.2. Then there exist constants $\lambda_0 \geq 0$ and N , depending only on d, m, p, δ, K , the function ω , such that*

$$\lambda \|\mathbf{u}\|_{L_p(\mathbb{R}^d)} + \sqrt{\lambda} \|D\mathbf{u}\|_{L_p(\mathbb{R}^d)} \leq N(\sqrt{\lambda} + 1) \|\mathcal{L}\mathbf{u} - \lambda\mathbf{u}\|_{W_p^{-1}(\mathbb{R}^d)} \quad (6)$$

holds for any $\mathbf{u} \in W_p^1(\mathbb{R}^d)$ and $\lambda \geq \lambda_0$. Moreover, for any $\mathbf{f}, \mathbf{g}_\alpha \in L_p(\mathbb{R}^d)$ and $\lambda > \lambda_0$, there exists a unique $\mathbf{u} \in W_p^1(\mathbb{R}^d)$ such that

$$\mathcal{L}\mathbf{u} - \lambda\mathbf{u} = \mathbf{f} + D_\alpha \mathbf{g}_\alpha \quad \text{in } \mathbb{R}^d.$$

3. PRELIMINARY RESULTS

Throughout this section we set

$$\bar{P}\mathbf{u} = -\mathbf{u}_t + A^{\alpha\beta} D_{\alpha\beta}\mathbf{u}, \quad (7)$$

where the entries of coefficient matrices $A^{\alpha\beta}$ are measurable functions of only $t \in \mathbb{R}$, i.e., $A_{ij}^{\alpha\beta} = A_{ij}^{\alpha\beta}(t)$, satisfying Assumption 2.1. With this operator \bar{P} we have the following theorem, which is proved in section 4, first for the case $p = 2$, and then for the general case $1 < p < \infty$.

Theorem 3.1. *Let $\mathbf{u} \in W_p^{1,2}(\mathbb{R}_T^{d+1})$, $T \in (-\infty, \infty]$. Then there exists a constant $N = N(d, m, p, \delta, K)$ such that*

$$\lambda \|\mathbf{u}\|_{L_p(\mathbb{R}_T^{d+1})} + \sqrt{\lambda} \|D\mathbf{u}\|_{L_p(\mathbb{R}_T^{d+1})} + \|D^2\mathbf{u}\|_{L_p(\mathbb{R}_T^{d+1})} + \|\mathbf{u}_t\|_{L_p(\mathbb{R}_T^{d+1})} \leq N \|\bar{P}\mathbf{u} - \lambda\mathbf{u}\|_{L_p(\mathbb{R}_T^{d+1})} \quad (8)$$

for all $\lambda \geq 0$. Moreover, for any $\lambda > 0$ and $\mathbf{f} \in L_p(\mathbb{R}_T^{d+1})$, there exists a unique $\mathbf{u} \in W_p^{1,2}(\mathbb{R}_T^{d+1})$ satisfying $\bar{P}\mathbf{u} - \lambda\mathbf{u} = \mathbf{f}$.

The above theorem, along with the arguments in [20], allows us to obtain the following lemmas and theorems.

Lemma 3.2. *Let $0 < r < R < \infty$ and $\mathbf{u} \in W_{p,loc}^{1,2}(\mathbb{R}^{d+1})$. Then*

$$\|\mathbf{u}_t\|_{L_p(Q_r)} + \|D^2\mathbf{u}\|_{L_p(Q_r)} \leq N (\|\bar{P}\mathbf{u}\|_{L_p(Q_R)} + \|D\mathbf{u}\|_{L_p(Q_R)} + \|\mathbf{u}\|_{L_p(Q_R)}),$$

where $N = N(d, m, p, \delta, K, r, R)$.

Proof. See the proof of Lemma 5.2 in [20]. \square

Lemma 3.3. *Let $0 < r < R < \infty$ and $\mathbf{u} \in C_{loc}^\infty(\mathbb{R}^{d+1})$. Assume that $\bar{P}\mathbf{u} = 0$ in Q_R . Then for any multi-index γ , we have*

$$\sup_{Q_r} |D^\gamma \mathbf{u}| + \sup_{Q_r} |D^\gamma \mathbf{u}_t| \leq N (\|D\mathbf{u}\|_{L_p(Q_R)} + \|\mathbf{u}\|_{L_p(Q_R)}),$$

where $N = N(|\gamma|, d, m, \delta, p, K, r, R)$.

Proof. See the proof of Lemma 5.8 in [20]. \square

Lemma 3.4. *Let $\mathbf{u} \in C_{loc}^\infty(\mathbb{R}^{d+1})$ and $\lambda \geq 0$. Assume that $\bar{P}\mathbf{u} - \lambda\mathbf{u} = 0$ in Q_2 . Then for any multi-index γ , we have*

$$\sup_{Q_1} |D^\gamma(D^2\mathbf{u})| + \sup_{Q_1} |D^\gamma\mathbf{u}_t| \leq N \left(\|\mathbf{u}_t\|_{L_p(Q_2)} + \|D^2\mathbf{u}\|_{L_p(Q_2)} + \sqrt{\lambda}\|D\mathbf{u}\|_{L_p(Q_2)} \right),$$

$$\sup_{Q_1} |D^\gamma(D\mathbf{u})| + \sup_{Q_1} |D^\gamma\mathbf{u}_t| \leq N \left(\|D\mathbf{u}\|_{L_p(Q_2)} + \sqrt{\lambda}\|\mathbf{u}\|_{L_p(Q_2)} \right),$$

where $N = N(|\gamma|, d, m, p, \delta, K)$.

Proof. See the proof of Lemma 5.9 and 7.3 in [20]. \square

Theorem 3.5. *Let $\lambda \geq 0$, $\kappa \geq 2$, and $r \in (0, \infty)$. Let $\mathbf{u} \in C_{loc}^\infty(\mathbb{R}^{d+1})$ such that $\bar{P}\mathbf{u} - \lambda\mathbf{u} = 0$ in $Q_{\kappa r}$. Then there is a constant N , depending only on d, m, p, δ , and K , such that*

$$\int_{Q_r} |D^2\mathbf{u}(t, x) - (D^2\mathbf{u})_{Q_r}|^p dx dt \leq N\kappa^{-p} \left(|D^2\mathbf{u}|^p + \lambda^{p/2}|D\mathbf{u}|^p \right)_{Q_{\kappa r}},$$

$$\int_{Q_r} |D\mathbf{u}(t, x) - (D\mathbf{u})_{Q_r}|^p dx dt \leq N\kappa^{-p} \left(|D\mathbf{u}|^p + \lambda^{p/2}|\mathbf{u}|^p \right)_{Q_{\kappa r}}.$$

Proof. See the proof of Theorem 5.10 and 7.4 in [20]. \square

By using the results above, one can obtain the following estimates, which are important in proving the main theorems. For their proofs, we refer to Theorem 5.1 and 7.1 in [20].

Theorem 3.6. *There is a constant $N = N(d, m, p, \delta, K)$ such that, for $\mathbf{u} \in W_{p,loc}^{1,2}(\mathbb{R}^{d+1})$, $r \in (0, \infty)$, and $\kappa \geq 4$,*

$$\int_{Q_r} |D^2\mathbf{u}(t, x) - (D^2\mathbf{u})_{Q_r}|^p dx dt \leq N\kappa^{d+2} (|\bar{P}\mathbf{u}|^p)_{Q_{\kappa r}} + N\kappa^{-p} (|D^2\mathbf{u}|^p)_{Q_{\kappa r}}. \quad (9)$$

Theorem 3.7. *Let $\mathbf{u} \in \mathcal{H}_{p,loc}^1(\mathbb{R}^{d+1})$, $\mathbf{g} = (\mathbf{g}_\alpha) \in L_{p,loc}(\mathbb{R}^{d+1})$, $r \in (0, \infty)$, and $\kappa \geq 4$. Assume $\bar{P}\mathbf{u} = D_\alpha\mathbf{g}_\alpha$ in $Q_{\kappa r}$. Then there is a constant $N = N(d, m, p, \delta, K)$ such that,*

$$\int_{Q_r} |D\mathbf{u}(t, x) - (D\mathbf{u})_{Q_r}|^p dx dt \leq N\kappa^{d+2} (|\mathbf{g}|^p)_{Q_{\kappa r}} + N\kappa^{-p} (|D\mathbf{u}|^p)_{Q_{\kappa r}}. \quad (10)$$

Remark 3.8. It is worth noting that Theorem 3.6 and 3.7 can be improved. Indeed, due to Lemma 6.2 and 8.2, following the proofs of Theorem 5.1 and 7.1 in [20] one can actually get

$$\int_{Q_r} |D^2\mathbf{u}(t, x) - (D^2\mathbf{u})_{Q_r}|^p dx dt \leq N\kappa^{d+2} (|\bar{P}\mathbf{u}|^p)_{Q_{\kappa r}} + N\kappa^{-p} (|D^2\mathbf{u}|^p)_{Q_{\kappa r}} \quad (11)$$

under the assumptions of Theorem 3.6, and

$$\int_{Q_r} |D\mathbf{u}(t, x) - (D\mathbf{u})_{Q_r}|^p dx dt \leq N\kappa^{d+2} (|\mathbf{g}|^p)_{Q_{\kappa r}} + N\kappa^{-p} (|D\mathbf{u}|^p)_{Q_{\kappa r}} \quad (12)$$

under the assumptions of Theorem 3.7.

4. SYSTEMS WITH COEFFICIENTS INDEPENDENT OF THE SPATIAL VARIABLES

In this section we prove Theorem 3.1, so recall that the operator \bar{P} was defined in (7), where $A^{\alpha\beta}$ are assumed to be matrices of measurable functions depending only on $t \in \mathbb{R}$, i.e., $A_{ij}^{\alpha\beta} = A_{ij}^{\alpha\beta}(t)$. Theorem 3.1, in case $p = 2$, is easily proved using Fourier transform methods.

Proof of Theorem 3.1 when $p = 2$. It is enough to prove the estimate (8) for $\lambda > 0$ and $\mathbf{u} \in C_c^\infty(\mathbb{R}^{d+1})$. Indeed, the estimate (8) for $\lambda = 0$ is proved by letting $\lambda \searrow 0$ once the estimate is established for $\lambda > 0$. To obtain the second assertion (solvability) of the theorem, we can just use the estimate (8), the solvability of the heat equation (i.e., $A_{ij}^{\alpha\beta} = \delta_{\alpha\beta}\delta_{ij}$), and the method of continuity.

For $\mathbf{u} \in C_c^\infty(\mathbb{R}^{d+1})$, set $\mathbf{f} = \bar{P}\mathbf{u} - \lambda\mathbf{u}$. Then by taking Fourier transforms in x -variables we see

$$-\tilde{\mathbf{u}}_t - \xi_\alpha \xi_\beta A^{\alpha\beta} \tilde{\mathbf{u}} - \lambda \tilde{\mathbf{u}} = \tilde{\mathbf{f}},$$

where $\tilde{\mathbf{u}} = (\tilde{u}^1, \dots, \tilde{u}^m)^T$ and $\tilde{u}^i(t, \xi)$, $i = 1, \dots, m$, are the Fourier transforms of u^i with respect to the spatial variables. By multiplying both sides of the above equation from the left by the complex conjugate of the Fourier transform of $(\lambda - \Delta)\mathbf{u}^T$, we obtain

$$\begin{aligned} -(\lambda + |\xi|^2) \bar{\tilde{\mathbf{u}}} \cdot \tilde{\mathbf{u}}_t - (\lambda + |\xi|^2) \xi_\alpha \xi_\beta \bar{\tilde{\mathbf{u}}}^T A^{\alpha\beta} \tilde{\mathbf{u}} - \lambda(\lambda + |\xi|^2) \bar{\tilde{\mathbf{u}}} \cdot \tilde{\mathbf{u}} \\ = (\lambda + |\xi|^2) \bar{\tilde{\mathbf{u}}} \cdot \tilde{\mathbf{f}} \end{aligned} \quad (13)$$

Note that

$$\begin{aligned} \Re(\bar{\tilde{\mathbf{u}}} \cdot \tilde{\mathbf{u}}_t) &= \frac{1}{2} \frac{\partial}{\partial t} |\tilde{\mathbf{u}}|^2, \quad \int_{-\infty}^T \frac{\partial}{\partial t} |\tilde{\mathbf{u}}(t, \xi)|^2 dt = |\tilde{\mathbf{u}}(T, \xi)|^2, \\ \Re(\xi_\alpha \xi_\beta \bar{\tilde{\mathbf{u}}}^T A^{\alpha\beta} \tilde{\mathbf{u}}) &= \xi_\alpha \xi_\beta \Re(\tilde{u}^i) A_{ij}^{\alpha\beta} \Re(\tilde{u}^j) + \xi_\alpha \xi_\beta \Im(\tilde{u}^i) A_{ij}^{\alpha\beta} \Im(\tilde{u}^j) \\ &\geq \delta |\xi|^2 |\tilde{\mathbf{u}}|^2. \end{aligned}$$

Thus by taking the real parts of the equation (13) and integrating them, we have

$$\delta \int_{\mathbb{R}_T^{d+1}} (\lambda + |\xi|^2)^2 |\tilde{\mathbf{u}}|^2 d\xi dt \leq - \int_{\mathbb{R}_T^{d+1}} (\lambda + |\xi|^2) \Re(\bar{\tilde{\mathbf{u}}} \cdot \tilde{\mathbf{f}}) d\xi dt. \quad (14)$$

We see that

$$-(\lambda + |\xi|^2) \Re(\bar{\tilde{\mathbf{u}}} \cdot \tilde{\mathbf{f}}) \leq (\lambda + |\xi|^2) \left(\varepsilon |\tilde{\mathbf{u}}|^2 + \frac{1}{4\varepsilon} |\tilde{\mathbf{f}}|^2 \right)$$

for any $\varepsilon > 0$. Set $\varepsilon = \delta(\lambda + |\xi|^2)/2$, then

$$\begin{aligned} - \int_{\mathbb{R}_T^{d+1}} (\lambda + |\xi|^2) \Re(\bar{\tilde{\mathbf{u}}} \cdot \tilde{\mathbf{f}}) d\xi dt &\leq \delta/2 \int_{\mathbb{R}_T^{d+1}} (\lambda + |\xi|^2)^2 |\tilde{\mathbf{u}}|^2 d\xi dt \\ &\quad + (2\delta)^{-1} \int_{\mathbb{R}_T^{d+1}} |\tilde{\mathbf{f}}|^2 d\xi dt. \end{aligned}$$

From this and (14) it follows that

$$\int_{\mathbb{R}_T^{d+1}} (\lambda + |\xi|^2)^2 |\tilde{\mathbf{u}}|^2 d\xi dt \leq \delta^{-2} \int_{\mathbb{R}_T^{d+1}} |\tilde{\mathbf{f}}|^2 d\xi dt.$$

This along with the Plancherel's theorem implies the inequality (8). \square

Let $\mathcal{Q} = \{Q_r(t, x) : (t, x) \in \mathbb{R}^{d+1}, r \in (0, \infty)\}$. For a function g defined on \mathbb{R}^{d+1} , we denote its (parabolic) maximal and sharp function, respectively, by

$$\begin{aligned} Mg(t, x) &= \sup_{Q \in \mathcal{Q}: (t, x) \in Q} \int_Q |g(s, y)| dy ds, \\ g^\#(t, x) &= \sup_{Q \in \mathcal{Q}: (t, x) \in Q} \int_Q |g(s, y) - (g)_Q| dy ds. \end{aligned}$$

Now that Theorem 3.1 is proved for the case $p = 2$, the results in section 3 are all available if $p = 2$. Then we proceed as follows to obtain the L_p -estimate, $1 < p < \infty$, for the parabolic system in (7).

Proposition 4.1. *There exists a constant $N = N(d, m, p, \delta, K)$ such that, for any $u \in C_c^\infty(\mathbb{R}^{d+1})$, we have*

$$\|D^2 \mathbf{u}\|_{L_p(\mathbb{R}^{d+1})} \leq N \|\bar{P} \mathbf{u}\|_{L_p(\mathbb{R}^{d+1})}.$$

Proof. Recall that the case $p = 2$ is proved above, so we first consider the case $p > 2$. From Theorem 3.6 along with an appropriate translation, we have

$$\begin{aligned} &(|D^2 \mathbf{u} - (D^2 \mathbf{u})_{Q_r(t_0, x_0)}|^2)_{Q_r(t_0, x_0)} \\ &\leq N \kappa^{d+2} (|\bar{P} \mathbf{u}|^2)_{Q_{\kappa r}(t_0, x_0)} + N \kappa^{-2} (|D^2 \mathbf{u}|^2)_{Q_{\kappa r}(t_0, x_0)} \end{aligned} \quad (15)$$

for $(t_0, x_0) \in \mathbb{R}^{d+1}$, $r > 0$ and $\kappa \geq 4$. Let

$$\mathcal{A}(t, x) = M(|\bar{P} \mathbf{u}|^2)(t, x), \quad \mathcal{B}(t, x) = M(|D^2 \mathbf{u}|^2)(t, x).$$

Then $(|\bar{P} \mathbf{u}|^2)_{Q_{\kappa r}(t_0, x_0)} \leq \mathcal{A}(t, x)$ for all $(t, x) \in Q_r(t_0, x_0)$. Similar inequalities are obtained for \mathcal{B} . From this and (15) it follows that, for any $(t, x) \in \mathbb{R}^{d+1}$ and $Q \in \mathcal{Q}$ such that $(t, x) \in Q$,

$$(|D^2 \mathbf{u} - (D^2 \mathbf{u})_Q|^2)_Q \leq N \kappa^{d+2} \mathcal{A}(t, x) + N \kappa^{-2} \mathcal{B}(t, x)$$

for $\kappa \geq 4$. Take the supremum of the left-hand side of the above inequality over all $Q \in \mathcal{Q}$ containing (t, x) . Also observe that

$$(|D^2 \mathbf{u} - (D^2 \mathbf{u})_Q|^2)_Q \leq (|D^2 \mathbf{u} - (D^2 \mathbf{u})_Q|^2)_Q.$$

Then we obtain

$$((D^2 \mathbf{u})^\#(t, x))^2 \leq N \kappa^{d+2} \mathcal{A}(t, x) + N \kappa^{-2} \mathcal{B}(t, x)$$

for $\kappa \geq 4$, where $N = N(d, m, \delta, K)$. Apply the Fefferman-Stein theorem on sharp functions and the Hardy-Littlewood maximal function theorem on the above inequality. More precisely,

$$\begin{aligned} \|D^2 \mathbf{u}\|_{L_p(\mathbb{R}^{d+1})} &\leq N \| (D^2 \mathbf{u})^\# \|_{L_p(\mathbb{R}^{d+1})} \\ &\leq N \kappa^{(d+2)/2} \|M(|\bar{P} \mathbf{u}|^2)\|_{L_{p/2}(\mathbb{R}^{d+1})}^{1/2} + N \kappa^{-1} \|M(|D^2 \mathbf{u}|^2)\|_{L_{p/2}(\mathbb{R}^{d+1})}^{1/2} \\ &\leq N \kappa^{(d+2)/2} \|\bar{P} \mathbf{u}\|_{L_p(\mathbb{R}^{d+1})} + N \kappa^{-1} \|D^2 \mathbf{u}\|_{L_p(\mathbb{R}^{d+1})}, \end{aligned}$$

where the last inequality is possible due to the assumption that $p > 2$. Now choose a large enough κ , then we see that the inequality in the theorem follows if $p > 2$. For the case $1 < p < 2$, we use the duality argument. \square

Let us complete the proof of Theorem 3.1 for all $1 < p < \infty$.

Proof of Theorem 3.1. First set $T = \infty$. As in the case $p = 2$, it is enough to prove the estimate in the theorem, which follows from the estimate proved in Proposition 4.1 and the Agmon's idea described in the proof of Theorem 4.1 in [19]. For the case $T < \infty$, we make use of the argument in the proof of Theorem 2.1 in [19], i.e. by using the fact that $\mathbf{u} = \mathbf{w}$ for $t < T$ where $\mathbf{w} \in W_p^{1,2}(\mathbb{R}^{d+1})$ solves $\bar{P}\mathbf{w} - \lambda\mathbf{w} = I_{t < T}(\bar{P}\mathbf{u} - \lambda\mathbf{u})$. The theorem is proved. \square

5. SYSTEMS IN NON-DIVERGENCE FORM WITH VMO_x COEFFICIENTS IN L_p

In this section we assume that the coefficients $A_{ij}^{\alpha\beta}(t, x)$ are measurable in $t \in \mathbb{R}$ and VMO in $x \in \mathbb{R}^d$. More precisely, Assumption 2.2 as well as Assumption 2.1 are satisfied.

Here is the main result of this section.

Theorem 5.1. *Let $\mathbf{u} \in W_p^{1,2}(\mathbb{R}_T^{d+1})$, $T \in (-\infty, \infty]$. Then there exist constants $\lambda_0 \geq 0$ and N , depending only on d, m, p, δ, K , and the function ω , such that*

$$\begin{aligned} \lambda \|\mathbf{u}\|_{L_p(\mathbb{R}_T^{d+1})} + \sqrt{\lambda} \|D\mathbf{u}\|_{L_p(\mathbb{R}_T^{d+1})} + \|D^2\mathbf{u}\|_{L_p(\mathbb{R}_T^{d+1})} \\ + \|\mathbf{u}_t\|_{L_p(\mathbb{R}_T^{d+1})} \leq N \|P\mathbf{u} - \lambda\mathbf{u}\|_{L_p(\mathbb{R}_T^{d+1})} \end{aligned}$$

for all $\lambda \geq \lambda_0$. Moreover, for any $\lambda > \lambda_0$ and $\mathbf{f} \in L_p(\mathbb{R}_T^{d+1})$, there exists a unique $\mathbf{u} \in W_p^{1,2}(\mathbb{R}_T^{d+1})$ satisfying $P\mathbf{u} - \lambda\mathbf{u} = \mathbf{f}$.

Remark 5.2. This theorem implies Theorem 2.3 for $p = q$. This is justified again by using the argument in the proof of Theorem 2.1 in [19].

To prove the above theorem, we need some preliminary results presented below. First recall that we have proved Theorem 3.1 in section 4, so the results in section 3 are available for all $p \in (1, \infty)$. Then we have

Lemma 5.3. *Let $B^\alpha = C = 0$. Also let $\mu, \nu \in (1, \infty)$, $1/\mu + 1/\nu = 1$, and $R \in (0, \infty)$. Then there exists a constant $N = N(d, m, q, \delta, K, \mu)$ such that, for any $\mathbf{u} \in C_c^\infty(\mathbb{R}^{d+1})$ vanishing outside Q_R , we have*

$$\begin{aligned} (|D^2\mathbf{u} - (D^2\mathbf{u})_{Q_r(t_0, x_0)}|^q)_{Q_r(t_0, x_0)} \leq N\kappa^{-q} (|D^2\mathbf{u}|^q)_{Q_{\kappa r}(t_0, x_0)} \\ + N\kappa^{d+2} \left((|P\mathbf{u}|^q)_{Q_{\kappa r}(t_0, x_0)} + \omega(R)^{1/\nu} (|D^2\mathbf{u}|^{q\mu})_{Q_{\kappa r}(t_0, x_0)}^{1/\mu} \right) \end{aligned}$$

for $r \in (0, \infty)$, $\kappa \geq 4$, and $(t_0, x_0) \in \mathbb{R}^{d+1}$.

Proof. Let $\kappa \geq 4$ and $r \in (0, \infty)$. We introduce another coefficients $\bar{A}^{\alpha\beta}$ defined as follows.

$$\begin{aligned} \bar{A}^{\alpha\beta}(t) &= \int_{B_{\kappa r}(t_0, x_0)} A^{\alpha\beta}(t, y) dy \quad \text{if } \kappa r < R, \\ \bar{A}^{\alpha\beta}(t) &= \int_{B_R} A^{\alpha\beta}(t, y) dy \quad \text{if } \kappa r \geq R. \end{aligned}$$

Set $\bar{P}\mathbf{u} = -\mathbf{u}_t + \bar{A}^{\alpha\beta} D_{\alpha\beta}\mathbf{u}$. Then from Theorem 3.6 (also using a translation) it follows that

$$\begin{aligned} (|D^2\mathbf{u} - (D^2\mathbf{u})_{Q_r(t_0, x_0)}|^q)_{Q_r(t_0, x_0)} \\ \leq N\kappa^{d+2} (|\bar{P}\mathbf{u}|^q)_{Q_{\kappa r}(t_0, x_0)} + N\kappa^{-q} (|D^2\mathbf{u}|^q)_{Q_{\kappa r}(t_0, x_0)}. \quad (16) \end{aligned}$$

Note that

$$\int_{Q_{\kappa r}(t_0, x_0)} |\bar{P}\mathbf{u}|^q dx dt \leq N \int_{Q_{\kappa r}(t_0, x_0)} |P\mathbf{u}|^q dx dt + N\chi, \quad (17)$$

where

$$\begin{aligned} \chi &= \int_{Q_{\kappa r}(t_0, x_0)} |(\bar{A}^{\alpha\beta} - A^{\alpha\beta})D_{\alpha\beta}\mathbf{u}|^q dx dt = \int_{Q_{\kappa r}(t_0, x_0) \cap Q_R} |(\bar{A}^{\alpha\beta} - A^{\alpha\beta})D_{\alpha\beta}\mathbf{u}|^q dx dt \\ &\leq \left(\int_{Q_{\kappa r}(t_0, x_0) \cap Q_R} |\bar{A} - A|^{q\nu} \right)^{1/\nu} \left(\int_{Q_{\kappa r}(t_0, x_0) \cap Q_R} |D^2\mathbf{u}|^{q\mu} \right)^{1/\mu} := I^{1/\nu} J^{1/\mu}, \end{aligned}$$

Using the definition of $\bar{A}^{\alpha\beta}$ and assumptions on $A^{\alpha\beta}$, we obtain the following estimates for I . If $\kappa r < R$,

$$\begin{aligned} I &\leq N \int_{t_0 - (\kappa r)^2}^{t_0} \int_{B_{\kappa r}(x_0)} |\bar{A} - A| dx dt \\ &\leq N(\kappa r)^{d+2} \omega(\kappa r) \leq N(\kappa r)^{d+2} \omega(R). \end{aligned}$$

In case $\kappa r \geq R$,

$$I \leq N \int_{-R^2}^0 \int_{B_R} |\bar{A} - A| dx dt \leq NR^{d+2} \omega(R) \leq N(\kappa r)^{d+2} \omega(R).$$

From the inequality (17) and the estimates for I , it follows that

$$(|\bar{P}\mathbf{u}|^q)_{Q_{\kappa r}(t_0, x_0)} \leq N\omega(R)^{1/\nu} (|D^2\mathbf{u}|^{q\mu})_{Q_{\kappa r}(t_0, x_0)}^{1/\mu} + N(|P\mathbf{u}|^q)_{Q_{\kappa r}(t_0, x_0)}.$$

This, together with (16), gives us

$$\begin{aligned} (|D^2\mathbf{u} - (D^2\mathbf{u})_{Q_r(t_0, x_0)}|^q)_{Q_r(t_0, x_0)} &\leq N\kappa^{d+2} \omega(R)^{1/\nu} (|D^2\mathbf{u}|^{q\mu})_{Q_{\kappa r}(t_0, x_0)}^{1/\mu} \\ &\quad + N\kappa^{d+2} (|P\mathbf{u}|^q)_{Q_{\kappa r}(t_0, x_0)} + N\kappa^{-q} (|D^2\mathbf{u}|^q)_{Q_{\kappa r}(t_0, x_0)} \end{aligned}$$

for any $r > 0$ and $\kappa \geq 4$. This finishes the proof. \square

Proposition 5.4. *Let $B^\alpha = C = 0$. Then there exist constants $R = R(d, m, p, \delta, K, \omega)$ and $N = N(d, m, p, \delta, K)$ such that, for $\mathbf{u} \in C_c^\infty(\mathbb{R}^{d+1})$ vanishing outside Q_R , we have*

$$\|D^2\mathbf{u}\|_{L_p(\mathbb{R}^{d+1})} \leq N\|P\mathbf{u}\|_{L_p(\mathbb{R}^{d+1})}.$$

Proof. Let \mathbf{u} be an infinitely differentiable function with compact support in Q_R , where R will be chosen below. Take $q > 1$ and $\mu > 1$ such that $1 < q\mu < p$. Let

$$\begin{aligned} \mathcal{A}(t, x) &= M(|P\mathbf{u}|^q)(t, x), \quad \mathcal{B}(t, x) = M(|D^2\mathbf{u}|^q)(t, x), \\ \mathcal{C}(t, x) &= (M(|D^2\mathbf{u}|^{q\mu})(t, x))^{1/\mu}. \end{aligned}$$

We note that $(|P\mathbf{u}|^q)_{Q_{\kappa r}(t_0, x_0)} \leq \mathcal{A}(t, x)$ for all $(t, x) \in Q_r(t_0, x_0)$. Similar inequalities hold true for \mathcal{B} and \mathcal{C} . Then by Lemma 5.3 it follows that (recall $1/\mu + 1/\nu = 1$)

$$(|D^2\mathbf{u} - (D^2\mathbf{u})_Q|^q)_Q \leq N\kappa^{-q}\mathcal{B}(t, x) + N\kappa^{d+2} \left(\mathcal{A}(t, x) + \omega(R)^{1/\nu}\mathcal{C}(t, x) \right)$$

for all $\kappa \geq 4$, $(t, x) \in \mathbb{R}^{d+1}$, and $Q \in \mathcal{Q}$ such that $(t, x) \in Q$. Take the supremum of the left-hand side of the above inequality over all $Q \in \mathcal{Q}$ containing (t, x) . Also observe that

$$(|D^2\mathbf{u} - (D^2\mathbf{u})_Q|^q)_Q \leq (|D^2\mathbf{u} - (D^2\mathbf{u})_Q|^q)_Q.$$

Then we obtain

$$\left((D^2 \mathbf{u})^\#(t, x) \right)^q \leq N \kappa^{d+2} \mathcal{A}(t, x) + N \kappa^{-q} \mathcal{B}(t, x) + N \kappa^{d+2} \omega(R)^{1/\nu} \mathcal{C}(t, x)$$

for all $\kappa \geq 4$, $(t, x) \in \mathbb{R}^{d+1}$, where $N = N(d, m, p, \delta, K)$. That is,

$$\begin{aligned} (D^2 \mathbf{u})^\#(t, x) &\leq N \kappa^{(d+2)/q} \mathcal{A}(t, x)^{1/q} + N \kappa^{-1} \mathcal{B}(t, x)^{1/q} \\ &\quad + N \kappa^{(d+2)/q} \omega(R)^{1/q\nu} \mathcal{C}(t, x)^{1/q}. \end{aligned}$$

Applying the Fefferman-Stein theorem on sharp functions and the Hardy-Littlewood maximal function theorem on the above inequality (recall $p > q\mu$), we have

$$\begin{aligned} \|D^2 \mathbf{u}\|_{L_p(\mathbb{R}^{d+1})} &\leq N \kappa^{(d+2)/q} \|\mathcal{A}^{1/q}\|_{L_p(\mathbb{R}^{d+1})} + N \kappa^{-1} \|\mathcal{B}^{1/q}\|_{L_p(\mathbb{R}^{d+1})} \\ &\quad + N \kappa^{(d+2)/q} \omega(R)^{1/(q\nu)} \|\mathcal{C}^{1/q}\|_{L_p(\mathbb{R}^{d+1})} \\ &\leq N \kappa^{(d+2)/q} \|P\mathbf{u}\|_{L_p(\mathbb{R}^{d+1})} + N \left(\kappa^{-1} + \kappa^{(d+2)/q} \omega(R)^{1/(q\nu)} \right) \|D^2 \mathbf{u}\|_{L_p(\mathbb{R}^{d+1})}, \end{aligned}$$

where $\kappa \geq 4$. Choose a big enough κ and then a small enough R , so that

$$N \left(\kappa^{-1} + \kappa^{(d+2)/q} \omega(R)^{1/(q\nu)} \right) < 1/2.$$

Then the estimate in the proposition follows. \square

Proof of Theorem 5.1. As noted earlier, it suffices to prove the estimate in the theorem. Thanks to Proposition 5.4, by using a partition of unity (see the proof of Theorem 5.7 in [19]), we obtain an estimate

$$\|D^2 \mathbf{u}\|_{L_p(\mathbb{R}^{d+1})} \leq N \left(\|P\mathbf{u}\|_{L_p(\mathbb{R}^{d+1})} + \|D\mathbf{u}\|_{L_p(\mathbb{R}^{d+1})} + \|\mathbf{u}\|_{L_p(\mathbb{R}^{d+1})} \right),$$

where $N = N(d, m, p, \delta, K, \omega)$ (this inequality is possible without having the condition $B^\alpha = C = 0$). Then using again the Agmon's idea in the proof of Theorem 4.1 in [19] and choosing a sufficiently large λ_0 , we arrive at the estimate in the theorem when $T = \infty$. To deal with the case $T < \infty$ we use again the argument in the proof of Theorem 2.1 in [19]. \square

6. PROOF OF THEOREM 2.3

Recall that by P we mean the operator defined in (1), where the matrices $A^{\alpha\beta}$, B^α , and C satisfy Assumption 2.1. Especially, the matrices $A^{\alpha\beta}$ satisfy Assumption 2.2.

The following two lemmas are possible since we have the L_p -estimate of systems with VMO_x coefficients (see Theorem 5.1). Their proofs can be done by imitating the proofs of Lemma 6.3 and Corollary 6.4 in [20].

Lemma 6.1. *Let $r \in (0, 1]$, $\kappa \in (1, \infty)$, and $\mathbf{u} \in W_{p,loc}^{1,2}(\mathbb{R}^{d+1})$. Then*

$$\|D^2 \mathbf{u}\|_{L_p(Q_r)} \leq N \left(\|P\mathbf{u}\|_{L_p(Q_{\kappa r})} + r^{-1} \|D\mathbf{u}\|_{L_p(Q_{\kappa r})} + r^{-2} \|\mathbf{u}\|_{L_p(Q_{\kappa r})} \right),$$

where N , independent of $r \in (0, 1]$, depends only on κ , d , m , p , δ , K and the function ω .

Lemma 6.2. *Let $p > q \geq 1$ and $r \in (0, 1]$. Let $B^\alpha = C = 0$ and assume that $\mathbf{u} \in W_{p,loc}^{1,2}(\mathbb{R}^{d+1})$ satisfies $P\mathbf{u} = 0$ in Q_{2r} . Then*

$$\left(|D^2 \mathbf{u}|^p \right)_{Q_r}^{1/p} \leq N \left(|D^2 \mathbf{u}| \right)_{Q_{2r}} \leq N \left(|D^2 \mathbf{u}|^q \right)_{Q_{2r}}^{1/q},$$

where N depends only on d , m , p , δ , K , and the function ω .

In the following, in order to prove Theorem 2.3, we shall use the idea in [20]. However, since we have parabolic *systems* and our statements are slightly different from those in [20], we present here some proofs.

Theorem 6.3. *Let $B^\alpha = C = 0$. Then there exists a constant $N = N(d, m, p, \delta, K, \omega)$ such that, for any $u \in C_c^\infty(\mathbb{R}^{d+1})$, $\kappa \geq 8$, and $r \in (0, 1/\kappa]$, we have*

$$\begin{aligned} \int_{Q_r} |D^2 \mathbf{u} - (D^2 \mathbf{u})_{Q_r}|^p dx dt \\ \leq N \kappa^{d+2} (|P\mathbf{u}|^p)_{Q_{\kappa r}} + N \left(\kappa^{-p} + \kappa^{d+2} \omega(\kappa r)^{1/2} \right) (|D^2 \mathbf{u}|^p)_{Q_{\kappa r}}. \end{aligned} \quad (18)$$

Proof. For given $\mathbf{u} \in C_c^\infty(\mathbb{R}^{d+1})$, $\kappa \geq 8$, and $r \in (0, 1/\kappa]$, find a unique function $\tilde{\mathbf{w}} \in W_p^{1,2}((-4, 3) \times \mathbb{R}^d)$ satisfying $\tilde{\mathbf{w}}(-4, \cdot) = 0$ and

$$P\tilde{\mathbf{w}} = \mathbf{f}I_{Q_{\kappa r}},$$

where $\mathbf{f} := P\mathbf{u}$. This is possible by Remark 5.2. In fact, $\tilde{\mathbf{w}} \in W_q^{1,2}((-4, 3) \times \mathbb{R}^d)$ for all $q \in (1, \infty)$ because $\mathbf{f}I_{Q_{\kappa r}} \in L_q((-4, 3) \times \mathbb{R}^d)$ for all $q \in (1, \infty)$. Let

$$\mathbf{w}(t, x) = \eta(t)\tilde{\mathbf{w}}(t, x),$$

where $\eta(t)$ is an infinitely differentiable function defined on \mathbb{R} such that

$$\eta(t) = 1, \quad -2 \leq t \leq 1, \quad \eta(t) = 0, \quad t \leq -3 \quad \text{or} \quad t \geq 2.$$

We see that $\mathbf{w} \in W_p^{1,2}(\mathbb{R}^{d+1})$ and, in addition, $\mathbf{w} \in W_q^{1,2}(\mathbb{R}^{d+1})$ for all $q \in (1, \infty)$. From Remark 5.2 (the estimate in Theorem 2.3 for $p = q$) we have

$$\int_{Q_{\kappa r}} |D^2 \mathbf{w}|^p dx dt \leq \int_{(-4, 3) \times \mathbb{R}^d} |D^2 \tilde{\mathbf{w}}|^p dx dt \leq N \int_{Q_{\kappa r}} |\mathbf{f}|^p dx dt,$$

where N depends only on d, m, p, δ, K , and the function ω . Thus

$$(|D^2 \mathbf{w}|^p)_{Q_{\kappa r}} \leq N (|\mathbf{f}|^p)_{Q_{\kappa r}}, \quad (19)$$

$$(|D^2 \mathbf{w}|^p)_{Q_r} \leq N \kappa^{d+2} (|\mathbf{f}|^p)_{Q_{\kappa r}}. \quad (20)$$

where $N = N(d, m, p, \delta, K, \omega)$.

Now we set

$$\mathbf{v} = \mathbf{u} - \mathbf{w}.$$

Then $\mathbf{v} \in W_p^{1,2}(\mathbb{R}^{d+1})$, $\mathbf{v} \in W_q^{1,2}(\mathbb{R}^{d+1})$, $q \in (1, \infty)$, and

$$P\mathbf{v} = 0 \quad \text{in} \quad Q_{\kappa r}.$$

Define \bar{P} by

$$\bar{P}\mathbf{u} = -\mathbf{u}_t + \bar{A}^{\alpha\beta}(t)D_{\alpha\beta}\mathbf{u},$$

where

$$\bar{A}^{\alpha\beta}(t) = \int_{B_{\kappa r/2}} A^{\alpha\beta}(t, y) dy.$$

Since $\mathbf{v} \in W_p^{1,2}(\mathbb{R}^{d+1})$ and $\kappa/2 \geq 4$, by Theorem 3.6 applied to the operator \bar{P} , we have

$$\int_{Q_r} |D^2 \mathbf{v}(t, x) - (D^2 \mathbf{v})_{Q_r}|^p dx dt \leq N \kappa^{d+2} (|\bar{P}\mathbf{v}|^p)_{Q_{\kappa r/2}} + N \kappa^{-p} (|D^2 \mathbf{v}|^p)_{Q_{\kappa r/2}}.$$

Using the fact that $P\mathbf{v} = 0$ in $Q_{\kappa r}$, we have

$$\begin{aligned} (|\bar{P}\mathbf{v}|^p)_{Q_{\kappa r/2}} &= (|\bar{P}\mathbf{v} - P\mathbf{v}|^p)_{Q_{\kappa r/2}} \\ &= \int_{Q_{\kappa r/2}} |(\bar{A}^{\alpha\beta}(t) - A^{\alpha\beta}(t, x)) D_{\alpha\beta}\mathbf{v}(t, x)|^p dx dt \\ &\leq N \left(\int_{Q_{\kappa r/2}} |\bar{A}(t) - A(t, x)|^{2p} dx dt \right)^{1/2} \left(\int_{Q_{\kappa r/2}} |D^2\mathbf{v}|^{2p} dx dt \right)^{1/2}, \end{aligned}$$

where we see

$$\int_{Q_{\kappa r/2}} |\bar{A}(t) - A(t, x)|^{2p} dx dt \leq N \int_{Q_{\kappa r/2}} |\bar{A}(t) - A(t, x)| dx dt \leq N\omega(\kappa r/2).$$

On the other hand, from Lemma 6.2 we see

$$\left(\int_{Q_{\kappa r/2}} |D^2\mathbf{v}|^{2p} dx dt \right)^{1/2} \leq N \left(\int_{Q_{\kappa r}} |D^2\mathbf{v}|^p dx dt \right),$$

where $N = N(d, m, p, \delta, K, \omega)$. Hence

$$\int_{Q_r} |D^2\mathbf{v}(t, x) - (D^2\mathbf{v})_{Q_r}|^p dx dt \leq N \left(\kappa^{-p} + \kappa^{d+2}\omega(\kappa r)^{1/2} \right) (|D^2\mathbf{v}|^p)_{Q_{\kappa r}}.$$

Note that

$$\begin{aligned} (|D^2\mathbf{v}|^p)_{Q_{\kappa r}} &\leq N (|D^2\mathbf{u}|^p)_{Q_{\kappa r}} + N (|D^2\mathbf{w}|^p)_{Q_{\kappa r}} \\ &\leq N (|D^2\mathbf{u}|^p)_{Q_{\kappa r}} + N (|\mathbf{f}|^p)_{Q_{\kappa r}}, \end{aligned}$$

where the second inequality is due to (19). Also note that, using the inequality (20), we have

$$\int_{Q_r} |D^2\mathbf{w}(t, x) - (D^2\mathbf{w})_{Q_r}|^p dx dt \leq N (|D^2\mathbf{w}|^p)_{Q_r} \leq N\kappa^{d+2} (|\mathbf{f}|^p)_{Q_{\kappa r}}.$$

Therefore,

$$\begin{aligned} &\int_{Q_r} |D^2\mathbf{u}(t, x) - (D^2\mathbf{u})_{Q_r}|^p dx dt \\ &\leq N \int_{Q_r} |D^2\mathbf{v}(t, x) - (D^2\mathbf{v})_{Q_r}|^p dx dt + N \int_{Q_r} |D^2\mathbf{w}(t, x) - (D^2\mathbf{w})_{Q_r}|^p dx dt \\ &\leq N \left(\kappa^{-p} + \kappa^{d+2}\omega(\kappa r)^{1/2} \right) (|D^2\mathbf{u}|^p + |\mathbf{f}|^p)_{Q_{\kappa r}} + N\kappa^{d+2} (|\mathbf{f}|^p)_{Q_{\kappa r}}. \end{aligned}$$

The theorem is proved. \square

Remark 6.4. As in Remark 3.8, one can replace the last term of (18) by

$$N \left(\kappa^{-p} + \kappa^{d+2}\omega(\kappa r)^{1/2} \right) (|D^2\mathbf{u}|^p)_{Q_{\kappa r}}.$$

If g is a function defined on \mathbb{R} , we define $(g)_{(a,b)}$ to be

$$(g)_{(a,b)} = \int_{(a,b)} g(s) ds = (b-a)^{-1} \int_a^b g(s) ds.$$

Especially, the maximal and sharp function of g are defined by

$$Mg(t) = \sup_{t \in (a,b)} \int_{(a,b)} |g(s)| ds,$$

$$g^\#(t) = \sup_{t \in (a,b)} \int_{(a,b)} |g(s) - (g)_{(a,b)}| ds,$$

where the supremums are taken over all intervals (a, b) containing t .

Using Theorem 6.3 and the argument in the proof of Corollary 3.2 in [20], we obtain the following corollary.

Corollary 6.5. *Let $B^\alpha = C = 0$. Then there exists a constant $N = N(d, m, p, \delta, K, \omega)$ such that, for any $\mathbf{u} \in C_c^\infty(\mathbb{R}^{d+1})$, $\kappa \geq 8$, and $r \in (0, 1/\kappa]$, we have*

$$\begin{aligned} & \int_{(-r^2, 0)} |\varphi(t) - (\varphi)_{(-r^2, 0)}|^p dt \\ & \leq N \kappa^{d+2} (\psi^p)_{(-(\kappa r)^2, 0)} + N \left(\kappa^{-p} + \kappa^{d+2} \omega(\kappa r)^{1/2} \right) (\varphi^p)_{(-(\kappa r)^2, 0)}, \end{aligned}$$

where

$$\varphi(t) = \|D^2 \mathbf{u}(t, \cdot)\|_{L_p(\mathbb{R}^d)}, \quad \psi(t) = \|P \mathbf{u}(t, \cdot)\|_{L_p(\mathbb{R}^d)}.$$

The result in the above corollary makes it possible to show that the sharp function of $\varphi(t)$ is pointwise bounded by the p -th root of the maximal functions of $|\psi|^p$ and $|\varphi|^p$ if $\varphi(t)$ has compact support.

Lemma 6.6. *Let $B^\alpha = C = 0$, $R \in (0, 1]$, and \mathbf{u} be a function in $C_c^\infty(\mathbb{R}^{d+1})$ such that $\mathbf{u}(t, x) = 0$ for $t \notin (0, R^4)$. Then*

$$\begin{aligned} \varphi^\#(t_0) & \leq N \kappa^{(d+2)/p} (M(\psi^p)(t_0))^{1/p} \\ & \quad + N \left((\kappa R)^{2-2/p} + \kappa^{-1} + \kappa^{(d+2)/p} (\omega(R))^{1/2p} \right) (M(\varphi^p)(t_0))^{1/p} \end{aligned}$$

for all $\kappa \geq 8$ and $t_0 \in \mathbb{R}$, where $N = N(d, m, p, \delta, K, \omega)$ and the functions φ and ψ are defined as in Corollary 6.5.

Proof. Fix κ such that $\kappa \geq 8$. If $r \leq R/\kappa$, then $\kappa r \leq R \leq 1$ and $A_{\kappa r}^\# \leq A_R^\# \leq \omega(R)$. Thus by Corollary 6.5,

$$\begin{aligned} & (|\varphi - (\varphi)_{(-r^2, 0)}|^p)_{(-r^2, 0)} \\ & \leq N \kappa^{d+2} (\psi^p)_{(-(\kappa r)^2, 0)} + N \left(\kappa^{-p} + \kappa^{d+2} (\omega(R))^{1/2} \right) (\varphi^p)_{(-(\kappa r)^2, 0)}. \end{aligned}$$

By an appropriate translation of this inequality we have

$$\begin{aligned} & \int_{(a,b)} |\varphi(t) - (\varphi)_{(a,b)}|^p dt \\ & \leq N \kappa^{d+2} (\psi^p)_{(c,b)} + N \left(\kappa^{-p} + \kappa^{d+2} (\omega(R))^{1/2} \right) (\varphi^p)_{(c,b)} \end{aligned}$$

if (a, b) is an interval such that $b - a \leq R^2/\kappa^2$ and $c = b - \kappa^2(b - a)$. Note that, for $t_0 \in (a, b)$,

$$(\psi^p)_{(c,b)} \leq M(\psi^p)(t_0), \quad (\varphi^p)_{(c,b)} \leq M(\varphi^p)(t_0).$$

Thus by the Hölder's inequality it follows that

$$\begin{aligned} \int_{(a,b)} |\varphi(t) - (\varphi)_{(a,b)}| &\leq N\kappa^{(d+2)/p} (M(\psi^p)(t_0))^{1/p} \\ &\quad + N \left(\kappa^{-1} + \kappa^{(d+2)/p} (\omega(R))^{1/2p} \right) (M(\varphi^p)(t_0))^{1/p}, \end{aligned}$$

where $t_0 \in (a, b)$ and $b-a \leq R^2/\kappa^2$. Now, if (a, b) is an interval such that $t_0 \in (a, b)$ and $b-a > R^2/\kappa^2$, then

$$\begin{aligned} \int_{(a,b)} |\varphi(t) - (\varphi)_{(a,b)}| dt &\leq 2 \int_{(a,b)} I_{(0,R^4)}(t) |\varphi(t)| dt \\ &\leq 2 \left(\int_{(a,b)} I_{(0,R^4)}(t) dt \right)^{1-1/p} \left(\int_{(a,b)} |\varphi(t)|^p dt \right)^{1/p} \\ &\leq 2 (\kappa R)^{2-2/p} (M(\varphi^p)(t_0))^{1/p}. \end{aligned}$$

Therefore, for all intervals $(a, b) \ni t_0$,

$$\begin{aligned} \int_{(a,b)} |\varphi(t) - (\varphi)_{(a,b)}| dt &\leq N\kappa^{(d+2)/p} (M(\psi^p)(t_0))^{1/p} \\ &\quad + N \left((\kappa R)^{2-2/p} + \kappa^{-1} + \kappa^{(d+2)/p} (\omega(R))^{1/2p} \right) (M(\varphi^p)(t_0))^{1/p}. \end{aligned}$$

Taking the supremum of the left-hand side of the above inequality over all intervals $(a, b) \ni t_0$, we obtain the inequality in the lemma. The lemma is proved. \square

We use again the Hardy-Littlewood theorem and Fefferman-Stein theorem to derive the $L_{q,p}$ -estimate of second order derivatives of solutions to parabolic systems.

Corollary 6.7. *Let $B^\alpha = C = 0$ and $1 < p < q < \infty$. Then there exists $R = R(d, m, p, \delta, K, \omega)$ such that, for any $\mathbf{u} \in C_c^\infty(\mathbb{R}^{d+1})$ satisfying $\mathbf{u}(t, x) = 0$ for $t \notin (0, R^4)$,*

$$\|D^2 \mathbf{u}\|_{L_{q,p}(\mathbb{R}^{d+1})} \leq N \|P\mathbf{u}\|_{L_{q,p}(\mathbb{R}^{d+1})},$$

where $N = N(d, m, p, q, \delta, K, \omega)$.

Proof. Let $\mathbf{u} \in C_c^\infty(\mathbb{R}^{d+1})$ be such that $\mathbf{u}(t, x) = 0$ for $t \notin (0, R^4)$, $R \in (0, 1]$, where R will be specified below. Using the inequality in Lemma 6.6 as well as the Hardy-Littlewood theorem and the Fefferman-Stein theorem (recall that $q/p > 1$), we arrive at

$$\begin{aligned} \|D^2 \mathbf{u}\|_{L_{q,p}(\mathbb{R}^{d+1})} &\leq N\kappa^{(d+2)/p} \|P\mathbf{u}\|_{L_{q,p}(\mathbb{R}^{d+1})} \\ &\quad + N \left((\kappa R)^{2-2/p} + \kappa^{-1} + \kappa^{(d+2)/p} (\omega(R))^{1/2p} \right) \|D^2 \mathbf{u}\|_{L_{q,p}(\mathbb{R}^{d+1})} \end{aligned}$$

for all $\kappa \geq 8$. Now we choose a large κ and then a small R such that

$$N \left((\kappa R)^{2-2/p} + \kappa^{-1} + \kappa^{(d+2)/p} (\omega(R))^{1/2p} \right) < \frac{1}{2}.$$

It then follows that

$$\|D^2 \mathbf{u}\|_{L_{q,p}(\mathbb{R}^{d+1})} \leq N \|P\mathbf{u}\|_{L_{q,p}(\mathbb{R}^{d+1})}.$$

This finishes the proof. \square

Proof of Theorem 2.3. For the case $p = q$, the theorem is proved by Theorem 5.1 (also see Remark 5.2). For the case $q > p$, the theorem is proved by the same reasoning in the proof of Theorem 5.1, i.e., using Corollary 6.7 and the arguments in [19]. \square

7. SYSTEMS IN DIVERGENCE FORM WITH VMO_x COEFFICIENTS IN L_p

This section is devoted to proving the following theorem, which is a counterpart of Theorem 5.1 for systems in the divergence form.

Theorem 7.1. *Let $\mathbf{u} \in \mathcal{H}_p^1(\mathbb{R}_T^{d+1})$, $T \in (-\infty, \infty]$. Then there exist constants $\lambda_0 \geq 0$ and N , depending only on d, m, p, δ, K , the function ω , such that*

$$\lambda \|\mathbf{u}\|_{L_p(\mathbb{R}_T^{d+1})} + \sqrt{\lambda} \|D\mathbf{u}\|_{L_p(\mathbb{R}_T^{d+1})} + \|\mathbf{u}_t\|_{\mathbb{H}_p^{-1}(\mathbb{R}_T^{d+1})} \leq N(\sqrt{\lambda} + 1) \|\mathcal{P}\mathbf{u} - \lambda\mathbf{u}\|_{\mathbb{H}_p^{-1}(\mathbb{R}_T^{d+1})}$$

for all $\lambda \geq \lambda_0$. Moreover, for any $\lambda > \lambda_0$ and $\mathbf{f}, \mathbf{g}_\alpha \in L_p(\mathbb{R}_T^{d+1})$, there exists a unique $\mathbf{u} \in \mathcal{H}_p^1(\mathbb{R}_T^{d+1})$ satisfying

$$\mathcal{P}\mathbf{u} - \lambda\mathbf{u} = \mathbf{f} + D_\alpha \mathbf{g}_\alpha.$$

Remark 7.2. As in Remark 5.2, Theorem 7.1 implies Theorem 2.4 for $p = q$.

For the proof of this theorem, we need a few auxiliary results. First, the following lemma is derived from Theorem 3.7 as Lemma 5.3 is derived from Theorem 3.6.

Lemma 7.3. *Let $B^\alpha = \hat{B}^\alpha = 0$, $C = 0$, $\mu, \nu \in (1, \infty)$, $1/\mu + 1/\nu = 1$, and $R \in (0, \infty)$. Assume $\mathbf{u} \in C_c^\infty(\mathbb{R}^{d+1})$ vanishing outside Q_R and $\mathcal{P}\mathbf{u} = D_\alpha \mathbf{g}_\alpha$, where $\mathbf{g} = (\mathbf{g}_\alpha) \in L_p(\mathbb{R}^{d+1})$. Then there exists a constant $N = N(d, m, q, \delta, K, \mu)$ such that*

$$\begin{aligned} (|D\mathbf{u} - (D\mathbf{u})_{Q_r(t_0, x_0)}|^q)_{Q_r(t_0, x_0)} &\leq N\kappa^{-q} (|D\mathbf{u}|^q)_{Q_{\kappa r}(t_0, x_0)} \\ &\quad + N\kappa^{d+2} \left((|\mathbf{g}|^q)_{Q_{\kappa r}(t_0, x_0)} + \omega(R)^{1/\nu} (|D\mathbf{u}|^{q\mu})_{Q_{\kappa r}(t_0, x_0)}^{1/\mu} \right) \end{aligned}$$

for any $r \in (0, \infty)$, $\kappa \geq 4$, and $(t_0, x_0) \in \mathbb{R}^{d+1}$.

Here is a counterpart of Proposition 5.4 which is proved by the same method.

Proposition 7.4. *Let $B^\alpha = \hat{B}^\alpha = 0$ and $C = 0$. Then there exist constants $R = R(d, m, p, \delta, K, \omega)$ and $N = N(d, m, p, \delta, K)$ such that, for $\mathbf{u} \in C_c^\infty(\mathbb{R}^{d+1})$ vanishing outside Q_R and $\mathbf{g} = (\mathbf{g}_\alpha) \in L_p(\mathbb{R}^{d+1})$ satisfying $\mathcal{P}\mathbf{u} = D_\alpha \mathbf{g}_\alpha$, we have*

$$\|D\mathbf{u}\|_{L_p(\mathbb{R}^{d+1})} \leq N\|\mathbf{g}\|_{L_p(\mathbb{R}^{d+1})}.$$

Corollary 7.5. *Let $\mathbf{f}, \mathbf{g} = (\mathbf{g}_\alpha) \in L_p(\mathbb{R}^{d+1})$ and $\mathbf{u} \in C_c^\infty(\mathbb{R}^{d+1})$. Then there exist R, λ_0 , and N , depending only on d, m, p, δ, K , and the function ω , such that if \mathbf{u} vanishes outside Q_R then we have*

$$\sqrt{\lambda} \|D\mathbf{u}\|_{L_p(\mathbb{R}^{d+1})} + \lambda \|\mathbf{u}\|_{L_p(\mathbb{R}^{d+1})} \leq N\sqrt{\lambda} \|\mathbf{g}\|_{L_p(\mathbb{R}^{d+1})} + N\|\mathbf{f}\|_{L_p(\mathbb{R}^{d+1})}$$

provided that $\lambda \geq \lambda_0$ and

$$\mathcal{P}\mathbf{u} - \lambda\mathbf{u} = D_\alpha \mathbf{g}_\alpha + \mathbf{f}.$$

Proof. Due to Proposition 7.4, the corollary can be proved by using the aforementioned idea of Agmon. See the proof of Lemma 5.5 of [19]. \square

Theorem 7.6. *Let $\mathbf{u} \in C_c^\infty(\mathbb{R}^{d+1})$. There exist λ_0 and N depending only on d, m, p, δ, K , and the function ω such that*

$$\begin{aligned} \|\mathbf{u}_t\|_{\mathbb{H}_p^{-1}(\mathbb{R}^{d+1})} + \sqrt{\lambda}\|D\mathbf{u}\|_{L_p(\mathbb{R}^{d+1})} + \lambda\|\mathbf{u}\|_{L_p(\mathbb{R}^{d+1})} \\ \leq N\sqrt{\lambda}\|\mathbf{g}\|_{L_p(\mathbb{R}^{d+1})} + N\|\mathbf{f}\|_{L_p(\mathbb{R}^{d+1})} \end{aligned}$$

provided that $\lambda \geq \lambda_0$ and

$$\mathcal{P}\mathbf{u} - \lambda\mathbf{u} = D_\alpha\mathbf{g}_\alpha + \mathbf{f}.$$

Moreover, there exists N depending only on d, m, p, δ, K , and ω such that

$$\begin{aligned} \|\mathbf{u}_t\|_{\mathbb{H}_p^{-1}(\mathbb{R}^{d+1})} + \sqrt{\lambda}\|D\mathbf{u}\|_{L_p(\mathbb{R}^{d+1})} + \lambda\|\mathbf{u}\|_{L_p(\mathbb{R}^{d+1})} \\ \leq N(\sqrt{\lambda} + 1)\|\mathcal{P}\mathbf{u} - \lambda\mathbf{u}\|_{\mathbb{H}_p^{-1}(\mathbb{R}^{d+1})} \quad (21) \end{aligned}$$

provided that $\lambda \geq \lambda_0$.

Proof. The first part of the theorem is obtained by noting that $(1 - \Delta)^{-1/2}$ and $(1 - \Delta)^{-1/2}D$ are bounded operators on $L_p(\mathbb{R}^{d+1})$ and using the standard partition of unity technique. Next we prove the second part. Denote

$$\mathbf{h} = \mathcal{P}\mathbf{u} - \lambda\mathbf{u}, \quad \bar{\mathbf{f}} = (1 - \Delta)^{-1}\mathbf{h}, \quad \bar{\mathbf{g}}_\alpha = -D_\alpha\bar{\mathbf{f}}.$$

Note that we have

$$D_\alpha\bar{\mathbf{g}}_\alpha + \bar{\mathbf{f}} = (1 - \Delta)\bar{\mathbf{f}} = \mathbf{h}.$$

Again due to the boundedness of $(1 - \Delta)^{-1/2}$ and $(1 - \Delta)^{-1/2}D_\alpha$ in $L_p(\mathbb{R}^{d+1})$, we have $\mathbf{h} \in \mathbb{H}_p^{-1}(\mathbb{R}^{d+1})$, $\bar{\mathbf{f}} \in L_p(\mathbb{R}^{d+1})$, $\bar{\mathbf{g}} = (\bar{\mathbf{g}}_\alpha) \in L_p(\mathbb{R}^{d+1})$ and

$$\|\bar{\mathbf{g}}\|_{L_p(\mathbb{R}^{d+1})} + \|\bar{\mathbf{f}}\|_{L_p(\mathbb{R}^{d+1})} \leq N\|\mathbf{h}\|_{\mathbb{H}_p^{-1}(\mathbb{R}^{d+1})}.$$

By the first part with $\bar{\mathbf{f}}$ and $\bar{\mathbf{g}}_\alpha$ in place of \mathbf{f} and \mathbf{g}_α , we bound the left-hand side of (21) by (recall $\mathbf{h} = D_\alpha\bar{\mathbf{g}}_\alpha + \bar{\mathbf{f}}$)

$$N\sqrt{\lambda}\|\bar{\mathbf{g}}\|_{L_p(\mathbb{R}^{d+1})} + N\|\bar{\mathbf{f}}\|_{L_p(\mathbb{R}^{d+1})} \leq N(\sqrt{\lambda} + 1)\|\mathbf{h}\|_{\mathbb{H}_p^{-1}(\mathbb{R}^{d+1})}.$$

The theorem is proved. \square

Proof of Theorem 7.1. As usual we only have to prove the apriori estimate. For $T = \infty$, this is given by Theorem 7.6. For $T < \infty$, we again make use of the argument in Theorem 2.1 in [20]. The theorem is proved. \square

8. PROOF OF THEOREM 2.4

Throughout this section, let \mathcal{P} be the operator defined in (2), where the matrices $A^{\alpha\beta}$, B^α , \hat{B}^α and C satisfy Assumption 2.1 and 2.2. We shall use the following Sobolev embedding type estimate.

Lemma 8.1. *Let $T \in (-\infty, \infty]$, $r \in (0, \infty)$, $1 < q \leq p < \infty$, and assume that*

$$1/q - 1/p \leq 1/(d+2).$$

Then for any function $\mathbf{u} \in \mathcal{H}_q^1(\mathbb{R}_T^{d+1})$ we have $\mathbf{u} \in L_p(\mathbb{R}_T^{d+1})$ and

$$\|\mathbf{u}\|_{L_p(\mathbb{R}_T^{d+1})} \leq N\|\mathbf{u}\|_{\mathcal{H}_q^1(\mathbb{R}_T^{d+1})},$$

where $N > 0$ depends only on d, m, q and p .

Proof. This result is implied by Theorem 7.1. See the proof of, for example, Lemma 8.1 of [20]. \square

With the aid of Lemma 8.1, we can get a counterpart of Lemma 6.2.

Lemma 8.2. *Let $r \in (0, 1]$, $q > 1$, and $B^\alpha = \hat{B}^\alpha = C = 0$. Assume that $\mathbf{u} \in \mathcal{H}_{q, \text{loc}}^1(\mathbb{R}^{d+1})$ and $\mathcal{P}\mathbf{u} = 0$ in Q_{2r} , then $D\mathbf{u} \in L_p(Q_r)$ and*

$$(|D\mathbf{u}|^p)_{Q_r}^{1/p} \leq N(|D\mathbf{u}|^q)_{Q_{2r}}^{1/q},$$

where N depends only on d, m, p, q, K, δ and ω .

Here comes the key estimates in proving Theorem 2.4. The first one is an analogue of Theorem 6.3, which is proved in a similar way. The second is a counterpart of Corollary 6.5.

Theorem 8.3. *Let $B^\alpha = \hat{B}^\alpha = 0$, $C = 0$, $\mathbf{g} = (\mathbf{g}_\alpha) \in L_{p, \text{loc}}(\mathbb{R}^{d+1})$, $\mathbf{u} \in \mathcal{H}_{p, \text{loc}}^1(\mathbb{R}^{d+1})$, $\kappa \geq 8$ and $r \in (0, 1/\kappa]$. Suppose $\mathcal{P}\mathbf{u} = D_\alpha \mathbf{g}_\alpha$. Then there exists a constant N depending only on $d, m, p, \delta, K, \omega$ such that*

$$\begin{aligned} \int_{Q_r} |D\mathbf{u} - (D\mathbf{u})_{Q_r}|^p dx dt \\ \leq N\kappa^{d+2}(|\mathbf{g}|^p)_{Q_{\kappa r}} + N\left(\kappa^{-p} + \kappa^{d+2}\omega(\kappa r)^{1/2}\right)(|D\mathbf{u}|^p)_{Q_{\kappa r}}. \end{aligned} \quad (22)$$

Proof. Without loss of generality we can assume that $\mathbf{u} \in \mathcal{H}_p^1(\mathbb{R}^{d+1})$. Due to Remark 7.2 there exists $\mathbf{v} \in \mathcal{H}_p^1((-4, 0) \times \mathbb{R}^d)$ satisfying

$$\mathcal{P}\mathbf{v} = D_\alpha(I_{Q_{\kappa r}} \mathbf{g}_\alpha).$$

Note that $\kappa r \leq 1$ so that $\mathbf{v}(t, x) = 0$ for $t \leq -1$. Remark 7.2 also gives that

$$\int_{(-4, 0) \times \mathbb{R}^d} |D\mathbf{v}|^p dx dt \leq N \int_{Q_{\kappa r}} |\mathbf{g}|^p dx dt. \quad (23)$$

Now we set $\mathbf{w} = \mathbf{u} - \mathbf{v}$ and notice that $\mathbf{w} \in \mathcal{H}_p^1((-4, 0) \times \mathbb{R}^d)$ and \mathbf{w} satisfies $\mathcal{P}\mathbf{w} = 0$ in $Q_{\kappa r}$. Define \bar{P} by

$$\bar{P}\mathbf{u} = -\mathbf{u}_t + \bar{A}^{\alpha\beta}(t)D_{\alpha\beta}\mathbf{u},$$

where

$$\bar{A}^{\alpha\beta}(t) = \int_{B_{\kappa r/2}} A^{\alpha\beta}(t, y) dy.$$

Then it is clear that in $Q_{\kappa r}$, we have

$$\bar{P}\mathbf{w} = D_\alpha \bar{\mathbf{g}}_\alpha, \quad \bar{\mathbf{g}}_\alpha = (\bar{A}^{\alpha\beta} - A^{\alpha\beta})D_\beta \mathbf{w}.$$

Applying Theorem 3.7 to \mathbf{w} instead of \mathbf{u} , using Hölder's inequality and Lemma 8.2, we obtain

$$\begin{aligned} \int_{Q_r} |D\mathbf{w}(t, x) - (D\mathbf{w})_{Q_r}|^p dx dt &\leq N\kappa^{d+2}(|\bar{\mathbf{g}}|^p)_{Q_{\kappa r/2}} + N\kappa^{-p}(|D\mathbf{w}|^p)_{Q_{\kappa r/2}} \\ &\leq N\kappa^{d+2}(|A - \bar{A}|^{2p})_{Q_{\kappa r/2}}^{1/2} (|D\mathbf{w}|^{2p})_{Q_{\kappa r/2}}^{1/2} + N\kappa^{-p}(|D\mathbf{w}|^p)_{Q_{\kappa r/2}} \\ &\leq N\kappa^{d+2}\omega(\kappa r)^{1/2}(|D\mathbf{w}|^p)_{Q_{\kappa r}} + N\kappa^{-p}(|D\mathbf{w}|^p)_{Q_{\kappa r/2}}. \end{aligned}$$

Observe that

$$(|D\mathbf{w}|^p)_{Q_{\kappa r/2}} \leq N(|D\mathbf{w}|^p)_{Q_{\kappa r}} \leq N(|D\mathbf{u}|^p)_{Q_{\kappa r}} + N(|D\mathbf{v}|^p)_{Q_{\kappa r}}.$$

These together with (23) imply that the left-hand side of (22) is bounded by

$$\begin{aligned} N \int_{Q_r} |D\mathbf{w} - (D\mathbf{w})_{Q_r}|^p dx dt + N \int_{Q_r} |D\mathbf{v}|^p dx dt \\ \leq N \left(\kappa^{-p} + \kappa^{d+2} \omega(\kappa r)^{1/2} \right) (|D\mathbf{u}|^p)_{Q_{\kappa r}} + N \kappa^{d+2} (|\mathbf{g}|^p)_{Q_{\kappa r}}. \end{aligned}$$

The theorem is proved. \square

Remark 8.4. As in Remark 3.8, one can replace the last term of (22) by

$$N \left(\kappa^{-p} + \kappa^{d+2} \omega(\kappa r)^{1/2} \right) (|D\mathbf{u}|^p)_{Q_{\kappa r}}.$$

Recall the notation $(g)_{(a,b)}$, $Mg(t)$ and $g^\#(t)$ defined in Section 6.

Corollary 8.5. Let $B^\alpha = \hat{B}^\alpha = 0$, $C = 0$. Assume $\mathbf{u} \in \mathcal{H}_p^1((S, T) \times \mathbb{R}^d)$, $\mathbf{g} = (\mathbf{g}_\alpha) \in L_p((S, T) \times \mathbb{R}^d)$ for some $-\infty < S < T < \infty$ and

$$\mathcal{P}\mathbf{u} = D_\alpha \mathbf{g}_\alpha \quad \text{in } (S, T) \times \mathbb{R}^d.$$

Then there exists a constant $N = N(d, m, p, \delta, K, \omega)$ such that, for any $\kappa \geq 8$ and $r \in (0, 1/\kappa]$, we have

$$\begin{aligned} \int_{(-r^2, 0)} |\varphi(t) - (\varphi)_{(-r^2, 0)}|^p dt \\ \leq N \kappa^{d+2} (\psi^p)_{(-(\kappa r)^2, 0)} + N \left(\kappa^{-p} + \kappa^{d+2} \omega(\kappa r)^{1/2} \right) (\varphi^p)_{(-(\kappa r)^2, 0)}, \end{aligned}$$

where

$$\varphi(t) = \|D\mathbf{u}(t, \cdot)\|_{L_p(\mathbb{R}^d)}, \quad \psi(t) = \|\mathbf{g}(t, \cdot)\|_{L_p(\mathbb{R}^d)}.$$

In the same fashion, we derive a counterpart of Lemma 6.6.

Lemma 8.6. Let $B^\alpha = \hat{B}^\alpha = 0$, $C = 0$, $R \in (0, 1]$, and $\mathbf{u} \in C_c^\infty(\mathbb{R}^{d+1})$, $\mathbf{g} = \mathbf{g}_\alpha \in L_p(\mathbb{R}^{d+1})$ such that $\mathbf{u}(t, x) = 0$ for $t \notin (0, R^4)$. Assume $\mathcal{P}\mathbf{u} = D_\alpha \mathbf{g}_\alpha$. Then

$$\begin{aligned} \varphi^\#(t_0) \leq N \kappa^{(d+2)/p} M(\psi^p)(t_0)^{1/p} \\ + N \left((\kappa R)^{2-2/p} + \kappa^{-1} + \kappa^{(d+2)/p} \omega(R)^{1/2p} \right) M(\varphi^p)(t_0)^{1/p} \end{aligned}$$

for all $\kappa \geq 8$ and $t_0 \in \mathbb{R}$, where $N = N(d, m, p, \delta, K, \omega)$ and the functions φ , ψ are defined as in Corollary 8.5.

Corollary 8.7. Let $B^\alpha = \hat{B}^\alpha = 0$, $C = 0$, $1 < p < q < \infty$ and $\mathbf{u} \in C_c^\infty(\mathbb{R}^{d+1})$, $\mathbf{g} = (\mathbf{g}_\alpha) \in L_{q,p}(\mathbb{R}^{d+1})$. Assume $\mathcal{P}\mathbf{u} = D_\alpha \mathbf{g}_\alpha$. Then there exists $R = R(d, m, p, \delta, K, \omega)$ such that, under the assumption $\mathbf{u}(t, x) = 0$ for $t \notin (0, R^4)$,

$$\|D\mathbf{u}\|_{L_{q,p}(\mathbb{R}^{d+1})} \leq N \|\mathbf{g}\|_{L_{q,p}(\mathbb{R}^{d+1})},$$

where $N = N(d, m, p, q, \delta, K, \omega)$.

Proof. It follows from Lemma 8.6, the Hardy-Littlewood theorem, and the Fefferman-Stein theorem. \square

By using a partition of unity, we further conclude the following corollary.

Corollary 8.8. *Let $B^\alpha = \hat{B}^\alpha = 0$, $C = 0$, $1 < p < q < \infty$ and $\mathbf{u} \in C_c^\infty(\mathbb{R}^{d+1})$, $\mathbf{g} = (\mathbf{g}_\alpha) \in L_{q,p}(\mathbb{R}^{d+1})$. Assume $\mathcal{P}\mathbf{u} = D_\alpha \mathbf{g}_\alpha$. Then we have,*

$$\|D\mathbf{u}\|_{L_{q,p}(\mathbb{R}^{d+1})} \leq N\|\mathbf{g}\|_{L_{q,p}(\mathbb{R}^{d+1})} + N\|\mathbf{u}\|_{L_{q,p}(\mathbb{R}^{d+1})},$$

where $N = N(d, m, p, q, \delta, K, \omega)$.

Proof of Theorem 2.4. As usual, it suffices to derive the apriori estimate. For the case $p = q$, the theorem follows from Theorem 7.1 and Remark 7.2. For the case $q > p$, the theorem is proved by again the same reasoning in the proof of Theorem 7.1, i.e., using Corollary 8.8, the Agmon's idea, and the arguments in [19]. Finally, in the case $q < p$, we use duality argument. \square

9. PROOF OF THEOREM 2.5 AND 2.6

In this section, we prove Theorem 2.5 and 2.6. Roughly speaking, the idea is that solutions of elliptic systems can be considered as steady state solutions of the corresponding parabolic systems. Therefore, the estimates of parabolic systems which we derived imply the estimates of elliptic systems.

Proof of Theorem 2.5. As usual, it suffices to prove (5). Let η be a smooth function on \mathbb{R} supported on $[-2, 2]$ and $\eta(t) = 1$ on $[-1, 1]$. For a fixed $T > 0$, denote $\mathbf{v}(t, x) = \eta(t/T)\mathbf{u}(x)$. Then it is clear that $\mathbf{v} \in W_p^{1,2}(\mathbb{R}^{d+1})$, and

$$(P - \lambda)\mathbf{v}(t, x) = -T^{-1}\eta'(t/T)\mathbf{u}(x) + \eta(t/T)(L - \lambda)\mathbf{u}(x). \quad (24)$$

Thanks to Theorem 5.1, we have

$$\lambda\|\mathbf{v}\|_{L_p(\mathbb{R}^{d+1})} + \sqrt{\lambda}\|D\mathbf{v}\|_{L_p(\mathbb{R}^{d+1})} + \|D^2\mathbf{v}\|_{L_p(\mathbb{R}^{d+1})} \leq N\|P\mathbf{v} - \lambda\mathbf{v}\|_{L_p(\mathbb{R}^{d+1})}.$$

This combined with (24) and the triangle inequality gives

$$\begin{aligned} & \|\eta(t/T)\|_{L_p(\mathbb{R})} \left(\lambda\|\mathbf{u}\|_{L_p(\mathbb{R}^d)} + \sqrt{\lambda}\|D\mathbf{u}\|_{L_p(\mathbb{R}^d)} + \|D^2\mathbf{u}\|_{L_p(\mathbb{R}^d)} \right) \\ & \leq N\|\eta(t/T)\|_{L_p(\mathbb{R})}\|L\mathbf{u} - \lambda\mathbf{u}\|_{L_p(\mathbb{R}^d)} + NT^{-1}\|\eta'(t/T)\|_{L_p(\mathbb{R}_T)}\|\mathbf{u}\|_{L_p(\mathbb{R}^d)}. \end{aligned}$$

Therefore,

$$\begin{aligned} & T^{1/p}\|\eta\|_{L_p(\mathbb{R})} \left(\lambda\|\mathbf{u}\|_{L_p(\mathbb{R}^d)} + \sqrt{\lambda}\|D\mathbf{u}\|_{L_p(\mathbb{R}^d)} + \|D^2\mathbf{u}\|_{L_p(\mathbb{R}^d)} \right) \\ & \leq NT^{1/p}\|\eta\|_{L_p(\mathbb{R})}\|L\mathbf{u} - \lambda\mathbf{u}\|_{L_p(\mathbb{R}^d)} + NT^{-1+1/p}\|\eta'\|_{L_p(\mathbb{R})}\|\mathbf{u}\|_{L_p(\mathbb{R}^d)}. \end{aligned}$$

Letting $T \rightarrow \infty$ yields (5). The theorem is proved. \square

Proof of Theorem 2.6. It is derived from Theorem 7.1 in the same way as Theorem 2.5 is derived from Theorem 5.1. We omit the details. \square

Remark 9.1. Finally, we remark that our main results can be extended to parabolic (or elliptic) systems of order $2n$, where n is any positive integer. The parabolic (or elliptic) condition in the sense of Petrovskii is assumed (see, for instance, [11], [12] and [22]). Naturally we require the leading coefficients to be VMO in x variables.

The details of the extension are omitted here and left to interested readers. We only point out that we should apply a generalized version of Lemma 5.4 in [20] involving higher order derivatives, and while using Agmon's idea instead of $\cos\sqrt{\lambda}y$ we consider the factor

$$e^{\lambda^{1/(2m)}\cos(\pi/(2m))y} \cos(\lambda^{1/(2m)}\sin(\pi/(2m))y),$$

which is a real solution of the ordinary differential equation

$$\frac{d^{2m}v}{dy^{2m}} = -\lambda v.$$

ACKNOWLEDGEMENT

The authors are grateful to Nicolai V. Krylov for very helpful discussions on the subject. They also thank Seick Kim for many useful comments on an earlier version of the manuscript.

REFERENCES

- [1] P. Acquistapace. On BMO regularity for linear elliptic systems. *Ann. Mat. Pura Appl.* (4) 161 (1992), 231–269.
- [2] P. Auscher. On necessary and sufficient conditions for L^p estimates of Riesz transforms associated to elliptic operators on \mathbb{R}^n and related estimates. *Mem. Amer. Math. Soc.*, 186 (2007), no. 871, xviii+75 pp.
- [3] S. Byun. $W^{1,p}$ regularity theory for parabolic equations in divergence form. *J. Evol. Equ.* 7 (2007), no. 3, 415–428.
- [4] S. Byun, H. Chen, M. Kim, L. Wang. L^p regularity theory for linear elliptic systems. *Discrete Contin. Dyn. Syst.* 18 (2007), no. 1, 121–134.
- [5] S. Campanato. Sistemi ellittici in forma divergenza. Regolarità all'interno. (Italian) [Elliptic systems in divergence form. Interior regularity] *Quaderni. [Publications] Scuola Normale Superiore Pisa, Pisa*, 1980. 187 pp.
- [6] S. Cho, H. Dong, S. Kim. On the fundamental solutions for strongly parabolic systems of second order. *Indiana Univ. Math. J.*, to appear.
- [7] F. Chiarenza, M. Frasca and P. Longo. Interior $W^{2,p}$ estimates for nondivergence elliptic equations with discontinuous coefficients. *Ricerche Mat.* 40 (1991) 149–168.
- [8] F. Chiarenza, M. Frasca and P. Longo. $W^{2,p}$ -solvability of the Dirichlet problem for nondivergence elliptic equations with VMO coefficients. *Trans. Amer. Math. Soc.* 336 (1993) 841–853.
- [9] R. Denk, M. Hieber and J. Prüss. R-boundedness, Fourier multipliers and problems of elliptic and parabolic type. *Mem. Amer. Math. Soc.* 166 (2003).
- [10] G. Di Fazio. L^p estimates for divergence form elliptic equations with discontinuous coefficients. *Boll. Un. Mat. Ital. A* (7) 10 (1996), no. 2, 409–420.
- [11] S.D. Eidel'man. *Parabolic Systems*. (1969) North-Holland, Amsterdam, London.
- [12] A. Friedman. *Partial Differential Equations of Parabolic Type*. (1964) Prentice-Hall, Englewood Cliffs, N.J.
- [13] Giaquinta, M. *Multiple integrals in the calculus of variations and nonlinear elliptic systems*. Princeton University Press:Princeton, NJ, 1983.
- [14] R. Haller-Dintelmann, H. Heck, M. Hieber. L^p - L^q -estimates for parabolic systems in non-divergence form with VMO coefficients. *J. London Math. Soc.*, (2) 74 (3) (2006), 717C736.
- [15] D. Kim. Parabolic equations with measurable coefficients, II. *J. Math. Anal. Appl.* 334 (2007), no. 1, 534–548.
- [16] D. Kim, N. V. Krylov. Elliptic differential equations with coefficients measurable with respect to one variable and VMO with respect to the others. *SIAM J. Math. Anal.*, 39 (2007), no. 2, 489–506/
- [17] D. Kim, N. V. Krylov. Parabolic equations with measurable coefficients. *Potential Anal.* 26 (2007), no. 4, 345–361.
- [18] N. V. Krylov. Parabolic equations in L^p -spaces with mixed norms. *Algebra i Analiz* 14 (4) (2002) 91–106 (in Russian); English translation: *St. Petersburg Math. J.* 14 (4) (2003) 603–614.
- [19] N. V. Krylov. Parabolic and elliptic equations with VMO coefficients. *Comm. Partial Differential Equations*, 32(3):453–475, (2007).
- [20] N. V. Krylov. Parabolic equations with VMO coefficients in spaces with mixed norms. *J. Funct. Anal.*, 250:521–558, (2007).
- [21] S. Leonardi and J. Kottas and J. Stara. Hölder regularity of the solutions of some classes of elliptic systems in convex nonsmooth domains. *Nonlinear Anal.*, 60 (2005), 925V944.

- [22] D. Palagachev, L. Softova. A priori estimates and precise regularity for parabolic systems with discontinuous data. *Discrete Contin. Dyn. Syst.* 13 (3) (2005) 721–742.
- [23] D. Palagachev, L. Softova. Characterization of the interior regularity for parabolic systems with discontinuous coefficients. *Atti. Accad. Naz. Lincei Cl. Sci. Fis. Mat. Natur. Rend. Lincei* (9) Mat. Appl. 16 (2005) 125–132.
- [24] D. Palagachev, L. Softova. Fine regularity for elliptic systems with discontinuous ingredients. *Arch. Math.* (Basel) 86 (2) (2006) 145–153.
- [25] M. A. Ragusa. Local Hölder regularity for solutions of elliptic systems. *Duke Math. J.* 113 (2002) 385–397.
- [26] V. A. Solonnikov. On boundary value problems for linear parabolic systems of differential equations of general form. (Russian) *Trudy Mat. Inst. Steklov.* 83 (1965) 3–163.
- [27] B. Stroffolini. Elliptic systems of PDE with BMO-coefficients. *Potential Anal.*, 15 (2001), 285–299.

(H. Dong) DIVISION OF APPLIED MATHEMATICS, BROWN UNIVERSITY, 182 GEORGE STREET,
PROVIDENCE, RI 02912, USA

E-mail address: `hdong@brown.edu`

(D. Kim) DEPARTMENT OF MATHEMATICS, UNIVERSITY OF SOUTHERN CALIFORNIA, 3620 SOUTH
VERMONT AVENUE, KAP 108, LOS ANGELES, CA 90089-2532, USA

E-mail address: `doyoonki@usc.edu`