

The Gauss-Seidel Numerical Procedure for Markov Stochastic Games

Harold J. Kushner*
Applied Mathematics Department
Lefschetz Center for Dynamical Systems
Brown University, Providence, RI 02912, USA
hjk@dam.brown.edu
401-863-1400

October 29, 2003

Abstract

Consider the problem of value iteration for solving Markov stochastic games. One simply iterates backwards, via a Jacobi-like procedure. The convergence of the Gauss-Seidel form of this procedure is shown for both the discounted and ergodic cost problems, under appropriate conditions, with extensions to problems where one stops when a boundary is hit or if any one of the players chooses to stop, with associated costs. Generally, the Gauss-Seidel procedure accelerates convergence.

Key words. Stochastic games, Markov games, Gauss-Seidel procedure, numerical algorithms

1 Introduction

We consider two-player, zero-sum, finite-state, Markov stochastic games. There are N states and, unless noted otherwise, we suppose that the controls are feedback and not randomized. In state i , player 1's (the minimizing player) control is denoted by u_i and that of player 2 (the maximizing player) is denoted by v_i . The convergence of the value iteration procedure (see (2.2) below) for Markov stochastic games for a discounted cost function (or where there is an absorbing boundary) was established in [7, 11]. The convergence of the Gauss-Seidel procedure was first established for the control problem in [10]. It is widely used and is generally faster than the Jacobi procedure; see, for example,

*This work was partially supported by Contract DAAD-19-02-1-0425 from the Army Research Office and National Science Foundation Grant ECS 0097447.

[5, 10, 9], where the role of the ordering of the states and preferred orderings are discussed. Indeed, it can be much faster—depending on the ordering of the states in the iteration. The references [5, 10] discuss the nature of the modified transition probability that Q represents (for the control problem) and shows why it is faster.

The convergence of the Gauss-Seidel form has not yet been established for the game problem. Under appropriate conditions, the convergence will be established for the discounted and ergodic cost functions, and for related problems such as where there is an absorbing boundary or optional stopping.

The u_i, v_i take values in compact sets that might depend on i . Define the control vectors $u = \{u_i, i \leq N\}$, $v = \{v_i, i \leq N\}$. Let $P(u, v) = \{p_{ij}(u_i, v_i); i, j \leq N\}$ denote the transition probabilities under controls u, v . For notational simplicity, it is assumed, when dealing with the discounted problem, that the discount factor $\rho \in (0, 1)$ is included in the $p_{ij}(u_i, v_i)$. Hence $P(u, v)$ is degenerate in that case: the row sums are $1 - \rho$. The cost rate when in state i and under u_i, v_i is the function $k_i(u_i, v_i)$. Let $\{X_n\}$ denote the random variables of chain. Then the discounted cost under u, v is

$$C_i(u, v) = E_i^{u, v} \sum_{n=1}^{\infty} \rho^n k_{X_n}(u_{X_n}, v_{X_n}),$$

where $E_i^{u, v}$ denotes the expectation under u, v and with initial state i . It is always supposed that the $p_{ij}(u_i, v_i)$ and $k_i(u_i, v_i)$ are continuous in the u_i, v_i . Define the vector $K(u, v) = \{k_i(u_i, v_i); i \leq N\}$.

In addition, unless noted otherwise, we assume that the Isaacs condition holds; namely, that for any N -vector $H = \{h_i, i \leq N\}$,

$$\sup_v \inf_u [P(u, v)H + K(u, v)] = \inf_u \sup_v [P(u, v)H + K(u, v)]. \quad (1.1)$$

In vector forms such as (1.1), it is always supposed that the inf and sup are taken line by line, so that the i th line is $\sup_{v_i} \inf_{u_i} [\sum_j p_{ij}(u_i, v_i)h_j + k_i(u_i, v_i)]$ and involves the inf and sup over u_i and v_i only. The condition (1.1) is used for notational simplicity. Otherwise, one must randomize the controls. Then, when the number of control values is finite, the control is replaced by the vector of probabilities, and the analog of (1.1) holds. If the controls take values in a continuum and (1.1) does not hold, then the randomization is more complicated, but the results can be readily extended. The condition (1.1) commonly holds for the games arising as numerical approximations to stochastic differential games under the conditions of [4, 6].

Section 2 concerns the discounted cost problem and also remarks on cases where there is forced stopping on hitting a boundary or with optional stopping. The ergodic cost problem is dealt with in Section 3 and the cost under u, v is

$$\gamma(u, v) = \lim_n \frac{1}{n} E_i^{u, v} \sum_{l=1}^n k_{X_l}(u_{X_l}, v_{X_l}). \quad (1.2)$$

In Section 3, it is first shown that the game version of a classical value iteration method converges. Then this is adapted to the Gauss-Seidel procedure. If controls \hat{u}^n, \hat{v}^n are used at time n , then write $p_{ij}^{(n)}(\hat{u}^n, \hat{v}^n; \hat{u}^{n-1}, \hat{v}^{n-1}; \dots; \hat{u}^1, \hat{v}^1)$ for the n -step transition probabilities. The ergodic cost problem uses the additional assumption that there is an $\epsilon > 0$, a state j_0 , and an integer $m \leq N$, such that

$$p_{ij_0}^{(m)}(\hat{u}^m, \hat{v}^m; \hat{u}^{m-1}, \hat{v}^{m-1}; \dots; \hat{u}^1, \hat{v}^1) \geq \epsilon \quad (1.3)$$

for all possible controls. This is a standard condition for the ergodic cost problem in the control literature [12]. See also [1, Vol 2] and [5, pp156–158]. Of particular interest are Markov chain games that arise as numerical approximations of games with diffusion models as in [4, 6], where (1.3) will commonly hold under the assumptions on the nondegeneracy of the diffusion in [4].

To date, there have not been proofs of the convergence of the Gauss-Seidel method for either the game or the control problem with ergodic cost criteria. Indeed, it does not always converge, even under (1.3) for ergodic models. But, it will converge if (1.3) holds for a modified transition probability. This will be discussed further in Section 3. The modified condition holds for the chains obtained as approximations in [4] under the nondegeneracy conditions used there. These chains are obtained via the Markov chain approximation methods of [9]. The book [2] discusses other numerical procedures, based on nonlinear programming methods. The paper [3] discusses what might be called a type of combined value iteration and approximation in policy space method.

2 The Discounted Cost Problem and Extensions

Until further notice, we consider the discounted cost case. Let \bar{C}_i denote the value of the game when starting in state i and define $\bar{C} = \{\bar{C}_i; i \leq N\}$. In vector form, the equation for the value is

$$\bar{C} = \sup_v \inf_u [P(u, v)\bar{C} + K(u, v)] = \inf_u \sup_v [P(u, v)\bar{C} + K(u, v)]. \quad (2.1)$$

In all such vector equations, The inf sup is taken by line; the i th line is over u_i, v_i . Recall that the discounting is incorporated into the $P(u, v)$. Hence, for any integer $m \geq 1$, $P^m(\hat{u}^m, \hat{v}^m; \dots; \hat{u}^1, \hat{v}^1)$ is a contraction (in the Euclidean norm sense) uniformly in the choices of the controls $\{\hat{u}^n, \hat{v}^n\}$. A unique solution \bar{C} exists and is the value [2, Theorem 3.1.1]. Let \bar{u}, \bar{v} denote any controls that realize (2.1).

Our aim is the computation of \bar{C} , hence of optimizing controls as well. A variety of computational methods are available. In [7, 8, 11] it was shown that, for any C^0 , the C^n in the iteration in value space algorithm

$$C^{n+1} = \sup_v \inf_u [P(u, v)C^n + K(u, v)] \quad (2.2)$$

converge to \bar{C} .

The Gauss-Seidel procedure for the game problem is the iteration in value space with successive substitutions, taken in the order $i = 1, 2, \dots$,

$$C_i^{n+1} = \sup_{v_i} \inf_{u_i} \left[\sum_{j=1}^{i-1} p_{ij}(u_i, v_i) C_j^{n+1} + \sum_{j=i}^N p_{ij}(u_i, v_i) C_j^n + k_i(u_i, v_i) \right]. \quad (2.3)$$

The convergence proof in Theorem 2.1 adapts the method of [5, 10]. The ordering of the states can vary with n .

Before proceeding, it is convenient to define a transition probability and cost vector that appears in the analysis. Consider the set of linear equations, where the vector C is given, solved by successive substitution in the order $i = 1, 2, \dots$:

$$D_i = \left[\sum_{j=1}^{i-1} p_{ij}(u_i, v_i) D_j + \sum_{j=i}^N p_{ij}(u_i, v_i) C_j + k_i(u_i, v_i) \right], \quad 1 \leq i \leq N. \quad (2.4)$$

This uniquely defines a matrix $Q(u, v) = \{q_{ij}(u, v); i, j \leq N\}$ and vector $\hat{K}(u, v) = \{\hat{k}_i(u, v); i \leq N\}$ such that $D = Q(u, v)C + \hat{K}(u, v)$. In detail, by successive substitutions in (2.4), we find that

$$\begin{aligned} q_{1j}(u, v) &= p_{1j}(u_1, v_1), \quad 1 \leq j \leq N, \\ q_{21}(u, v) &= p_{21}(u_2, v_2) q_{11}(u, v), \\ q_{2j}(u, v) &= p_{2j}(u_2, v_2) + p_{21}(u_2, v_2) q_{1j}(u, v), \quad 2 \leq j \leq N. \end{aligned}$$

In general,

$$\begin{aligned} q_{ij}(u, v) &= p_{ij}(u_i, v_i) + \sum_{k=1}^{i-1} p_{ik}(u_i, v_i) q_{kj}(u, v), \quad j \geq i, \\ q_{ij}(u, v) &= \sum_{k=1}^{i-1} p_{ik}(u_i, v_i) q_{kj}(u, v), \quad 1 \leq j < i. \end{aligned} \quad (2.5)$$

$Q(u, v)$ can also be defined from (2.4) in terms of the upper and lower triangular matrices formed from $P(u, v)$, but we prefer to write the details. Also,

$$\begin{aligned} \hat{k}_1(u, v) &= k_1(u_1, v_1), \\ \hat{k}_2(u, v) &= p_{21}(u_2, v_2) \hat{k}_1(u, v) + k_2(u_2, v_2), \end{aligned}$$

and, in general,

$$\hat{k}_i(u, v) = \sum_{k=1}^{i-1} p_{ik}(u_i, v_i) \hat{k}_k(u, v) + k_i(u_i, v_i). \quad (2.6)$$

Note that, for the discounted cost problem where the discount factor is incorporated into the $p_{ij}(u_i, v_i)$, $Q(u, v)$ is a degenerate transition matrix since the row sums satisfy $\sum_j q_{ij}(u, v) \leq \rho$ for all i and controls. If there is no discounting,

then the row sums are always unity. These facts are easily proved by induction, starting with $i = 1$.

Theorem 2.1. *For any C^0 , the C^n in (2.3) converges to \bar{C} .*

Proof. Since

$$\bar{C}_i = \sup_{v_i} \inf_{u_i} \left[\sum_{j=1}^{i-1} p_{ij}(u_i, v_i) \bar{C}_j + \sum_{j=i}^N p_{ij}(u_i, v_i) \bar{C}_j + k_i(u_i, v_i) \right], \quad (2.7)$$

by successive substitutions, we can write (2.1) in the equivalent form

$$\bar{C} = \sup_v \inf_u \left[Q(u, v) \bar{C} + \hat{K}(u, v) \right] = Q(\bar{u}, \bar{v}) \bar{C} + \hat{K}(\bar{u}, \bar{v}). \quad (2.8)$$

Similarly, with u^n, v^n realizing (2.3), the following is equivalent to (2.3):

$$C^{n+1} = \sup_v \inf_u \left[Q(u, v) C^n + \hat{K}(u, v) \right] = [Q(u^n, v^n) C^n + \hat{K}(u^n, v^n)]. \quad (2.9)$$

In (2.8) and (2.9), it is understood that the inf and sup are again taken line by line, in the order $i = 1, 2, \dots$. The inf and sup in line 1 are over u_1 and v_1 , and in turn, that in line i are over u_i, v_i .

For any u, v , and $i = 1, \dots, N$, (2.1) yields

$$\begin{aligned} & \sum_{j=1}^{i-1} p_{ij}(\bar{u}_i, v_i) \bar{C}_j + \sum_{j=i}^N p_{ij}(\bar{u}_i, v_i) \bar{C}_j + k_i(\bar{u}_i, v_i) \\ & \leq \bar{C}_i = \sup_{v_i} \inf_{u_i} \left[\sum_{j=1}^{i-1} p_{ij}(u_i, v_i) \bar{C}_j + \sum_{j=i}^N p_{ij}(u_i, v_i) \bar{C}_j + k_i(u_i, v_i) \right] \\ & = \sum_{j=1}^{i-1} p_{ij}(\bar{u}_i, \bar{v}_i) \bar{C}_j + \sum_{j=i}^N p_{ij}(\bar{u}_i, \bar{v}_i) \bar{C}_j + k_i(\bar{u}_i, \bar{v}_i) \\ & \leq \sum_{j=1}^{i-1} p_{ij}(u_i, \bar{v}_i) \bar{C}_j + \sum_{j=i}^N p_{ij}(u_i, \bar{v}_i) \bar{C}_j + k_i(u_i, \bar{v}_i). \end{aligned} \quad (2.10)$$

In vector notation, this can be written as

$$P(\bar{u}, v) \bar{C} + K(\bar{u}, v) \leq \bar{C} \leq P(u, \bar{v}) \bar{C} + K(u, \bar{v}).$$

It can also be written as

$$Q(\bar{u}, v) \bar{C} + \hat{K}(\bar{u}, v) \leq \bar{C} \leq Q(u, \bar{v}) \bar{C} + \hat{K}(u, \bar{v}). \quad (2.11)$$

For any u, v , (2.3) or, equivalently, (2.9) yields

$$\begin{aligned} & Q(u^n, v) C^n + \hat{K}(u^n, v) \\ & \leq C^{n+1} = \sup_v \inf_u \left[Q(u, v) C^n + \hat{K}(u, v) \right] \leq Q(u, v^n) C^n + \hat{K}(u, v^n). \end{aligned} \quad (2.12)$$

Selecting $(u, v) = (u^n, v^n)$ in (2.11) and $(u, v) = (\bar{u}, \bar{v})$ in (2.12) yields

$$\begin{aligned}
Q(u^n, \bar{v})(C^{n+1} - \bar{C}) &= \left[Q(u^n, \bar{v})C^{n+1} + \hat{K}(u^n, \bar{v}) \right] - \left[Q(u^n, \bar{v})\bar{C} + \hat{K}(u^n, \bar{v}) \right] \\
&\leq C^{n+1} - \bar{C} \\
&\leq \left[Q(\bar{u}, v^n)C^n + \hat{K}(\bar{u}, v^n) \right] - \left[Q(\bar{u}, v^n)\bar{C} + \hat{K}(\bar{u}, v^n) \right] \\
&= Q(\bar{u}, v^n)(C^n - \bar{C}).
\end{aligned} \tag{2.13}$$

Iterating yields

$$\left[Q(u^n, \bar{v}) \cdots Q(u^1, \bar{v}) \right] (C^1 - \bar{C}) \leq C^{n+1} - \bar{C} \leq \left[Q(\bar{u}, v^n) \cdots Q(\bar{u}, v^1) \right] (C^1 - \bar{C}). \tag{2.14}$$

For the discounted problem the row sums of $Q(u, v)$ satisfy $\sum_j q_{ij}(u, v) \leq \rho$ for all i and controls. Hence $C^n \rightarrow \bar{C}$ as $n \rightarrow \infty$. ■

Stopping when hitting a boundary set. Now, we allow $\rho \in (0, 1]$, so that the discounting can be dropped if desired. Suppose that the process stops when a boundary set is hit and that the mean time to reach the boundary set is bounded, uniformly in the controls and initial condition. Thus we can suppose that the boundary set is absorbing and has zero cost. Without loss of generality, let 0 denote the boundary state. Let $P(u, v)$ still denote the matrix of transition probabilities among the states $1, \dots, N$ only. With $\bar{C} = \{\bar{C}_i; 1 \leq i \leq N\}$ and $C^n = \{C_i^n; 1 \leq i \leq N\}$ the products on either side of (2.14) go to zero. Thus $C^n \rightarrow \bar{C}$. It is preferable if state 1 is connected to the boundary and the states are ordered so that the “mean flow” is toward the boundary as one goes from the lower to the higher numbered states, where possible.

Optional stopping problems. As in the above paragraph, let $\rho \in (0, 1]$. Various forms of optional stopping can be handled. There are now three ways that the process can be stopped. One is by hitting a predefined stopping set, denoted by state 0, as in the previous paragraph. Call the time τ_0 . Otherwise, either player can decide to have the game stopped. The associated times are called τ_i for player i . After stopping for whatever reason, the state goes to absorbing 0, with zero holding cost there. The $P(u, v)$ represents the transition probabilities only among the states $1, \dots, N$. For given functions $g_i(\cdot)$, the cost is now

$$\begin{aligned}
C_i(u, v) &= E_i^{u, v} \sum_{n=0}^{\tau_0 \wedge \tau_1 \wedge \tau_2 - 1} k_{X_n}(u_{X_n}, v_{X_n}) \\
&\quad + E_i^{u, v} g_1(X_{\tau_1}) I_{\{\tau_1 \leq \tau_2, \tau_1 < \tau_0\}} + E_i^{u, v} g_2(X_{\tau_2}) I_{\{\tau_2 < \tau_1 \wedge \tau_0\}}.
\end{aligned} \tag{2.15}$$

The controls u_i, v_i can now take the new value *stop* as well as the original values used in Theorem 2.1. Let $k_i(u_i, v_i) \geq \epsilon > 0$ for all i, u_i, v_i values other than the value *stop*, and suppose that $g_1(\cdot) \neq g_2(\cdot)$ but $g_1(i) \geq g_2(i)$. Extend the definition of the $k_i(\cdot)$ to include the control value *stop*, by writing $k_i(\text{stop}, v_i) = g_1(i)$ and let $k_i(u_i, \text{stop}) = g_2(i)$ if $u_i \neq \text{stop}$ and let it be zero otherwise. Then

the Gauss-Seidel algorithm can be written as (2.3). We have $g_2(i) \leq C_i^n \leq g_1(i)$. Similarly (2.1) holds and $g_2(i) \leq \bar{C}_i \leq g_1(i)$.

Let (u^n, v^n) satisfy (2.3) and (\bar{u}, \bar{v}) satisfy (2.1). Due to the positivity of $k_i(u_i, v_i)$, for u_i and/or v_i not equal to *stop*, if player 1 uses u^n and player 2 uses some \hat{v}^n at time n , then $P(u^n, \hat{v}^n) \cdots P(u^0, \hat{v}^0) \rightarrow 0$ and $Q(u^n, \hat{v}^n) \cdots Q(u^0, \hat{v}^0) \rightarrow 0$ as $n \rightarrow \infty$ uniformly in the $\{\hat{v}^n\}$ choices. Analogously, $Q(\bar{u}, \hat{v}^n) \cdots Q(\bar{u}, \hat{v}^0) \rightarrow 0$ for all $\{\hat{v}^n\}$ choices. Using these facts and following the logic of the proof of Theorem 2.1 yields the convergence $C^n \rightarrow \bar{C}$ for this problem.

3 The Ergodic Cost Problem

Now $P(u, v)$ is the transition matrix for a controlled Markov chain which is ergodic under any u, v . We adapt the procedure of [5, pp156–158], originally due to White [12]. Let \mathbf{e} denote the N -vector, all of whose components are unity.

A Jacobi procedure. We first consider the analog of the simple backwards iteration (Jacobi) procedure (2.2), whose convergence for the game with ergodic payoffs has not been proved to date in the literature. For arbitrary W^0 , define the vectors W^n, w^n recursively by

$$\begin{aligned} W^n &= \sup_v \inf_u [P(u, v)w^{n-1} + K(u, v)] \\ w^n &= W^n - W_{j_0}^n \mathbf{e}, \end{aligned} \tag{3.1}$$

where j_0 is defined above (1.3). There is a value for the game [2, Section 5.2]. The value $\bar{\gamma}$ is given by

$$\bar{W} + \bar{\gamma} \mathbf{e} = \sup_v \inf_u [P(u, v)\bar{W} + K(u, v)]. \tag{3.2}$$

As for the control problem, the value of \bar{W} is unique, up to the addition of a vector with constant components, and the value of $\bar{\gamma}$ is unique. An alternative way of writing (3.2) is as

$$\begin{aligned} \bar{W} &= \sup_v \inf_u [P(u, v)\bar{w} + K(u, v)] \\ \bar{w} &= \bar{W} - \bar{W}_{j_0} \mathbf{e}. \end{aligned} \tag{3.2a}$$

Theorem 3.1. *w^n converges to the value $\bar{\gamma}$ of the game.*

Proof. Recall the condition (1.3) and the definitions of m and j_0 there. Let u^n, v^n be the selected controls in (3.1). Define $c_n = W_{j_0}^n \mathbf{e}$. Then, for any u, v ,

$$\begin{aligned} &P(u^n, v)w^{n-1} + K(u^n, v) \\ &\leq W^n = P(u^n, v^n)w^{n-1} + K(u^n, v^n) \leq P(u, v^n)w^{n-1} + K(u, v^n). \end{aligned} \tag{3.3}$$

Let $(u, v) = (u^{n-1}, v^{n-1})$ in (3.3) and use the definition of w^n in (3.1) to get

$$\begin{aligned} & P(u^n, v^{n-1})w^{n-1} + K(u^n, v^{n-1}) - c_n \\ & \leq w^n \leq P(u^{n-1}, v^n)w^{n-1} + K(u^{n-1}, v^n) - c_n. \end{aligned}$$

Replacing n with $n - 1$ in (3.3) and letting $(u, v) = (u^n, v^n)$ yields, for $i \leq N$,

$$\begin{aligned} & P(u^{n-1}, v^n)w^{n-2} + K(u^{n-1}, v^n) - c_{n-1} \\ & \leq w^{n-1} \leq P(u^n, v^{n-1})w^{n-2} + K(u^n, v^{n-1}) - c_{n-1}. \end{aligned}$$

The last two inequalities yield

$$\begin{aligned} & P(u^n, v^{n-1})(w^{n-1} - w^{n-2}) - (c_n - c_{n-1}) \leq w^n - w^{n-1} \\ & \leq P(u^{n-1}, v^n)(w^{n-1} - w^{n-2}) - (c_n - c_{n-1}). \end{aligned} \quad (3.4)$$

Iterating (3.4) $m - 1$ times leads to

$$\begin{aligned} & P(u^n, v^{n-1}) \cdots P(u^{n-m+1}, v^{n-m})(w^{n-m} - w^{n-m-1}) - (c_n - c_{n-m}) \\ & \leq w^n - w^{n-1} \\ & \leq P(u^{n-1}, v^n) \cdots P(u^{n-m}, v^{n-m+1})(w^{n-m} - w^{n-m-1}) - (c_n - c_{n-m}). \end{aligned} \quad (3.5)$$

Define $\delta w^n = w^n - w^{n-1} = \{\delta w_i^n; i \leq N\}$. Then the right hand inequality of (3.5) yields, for $i \leq N$,

$$\begin{aligned} & \delta w_i^n \leq \\ & \sum_j p_{ij}(u^{n-1}, v^n; u^{n-2}, v^{n-1}; \dots; u^{n-m}, v^{n-m+1}) \delta w_j^{n-m} - [W_{j_0}^n - W_{j_0}^{n-m}]. \end{aligned} \quad (3.6)$$

Since $w_{j_0}^n = 0$ for all n , we have $\delta w_{j_0}^n = 0$. This, with (3.6) and (from (1.3))

$$p_{ij_0}^{(m)}(u^{n-1}, v^n; u^{n-2}, v^{n-1}; \dots; u^{n-m}, v^{n-m+1}) \geq \epsilon > 0$$

for all i, n , and controls, yields for $i \leq N$

$$\max_i \delta w_i^n \leq (1 - \epsilon) \max_j \delta w_j^{n-m} - [W_{j_0}^n - W_{j_0}^{n-m}].$$

Analogously, using the fact that $\min_i w_i^n \leq 0$ and that

$$p_{ij_0}(u^n, v^{n-1}; u^{n-1}, v^{n-2}; \dots; u^{n-m+1}, v^{n-m}) \geq \epsilon$$

for all i, n , the left hand inequality of (3.5) yields, for $i \leq N$,

$$\min_i \delta w_i^n \geq (1 - \epsilon) \min_j \delta w_j^{n-m} - [W_{j_0}^n - W_{j_0}^{n-m}].$$

Hence, for all i ,

$$\left[\max_i - \min_i \right] \delta w_i^n \leq (1 - \epsilon) \left[\max_i - \min_i \right] \delta w_i^{n-m}, \quad (3.7)$$

which implies that w^n converges to, say, \bar{w} . Hence W^n converges to, say, \bar{W} , and the limits satisfy (3.2a). Hence, (3.2) holds with $\bar{\gamma} = \bar{w}_{j_0}$. ■

The Gauss-Seidel procedure. The Gauss-Seidel form of (3.1) is, in order $i = 1, 2, \dots$,

$$W_i^n = \sup_{v_i} \inf_{u_i} \left[\sum_{j=1}^{i-1} p_{ij}(u_i, v_i) W_j^n + \sum_{j=i}^N p_{ij}(u_i, v_i) [W_j^{n-1} - W_{j_0}^{n-1}] + k_i(u_i, v_i) \right], \quad (3.8)$$

Recall the definition of $Q(u, v)$ and $\hat{K}(u, v)$ from Section 2. Then, in matrix notation, (3.8) can be written as

$$\begin{aligned} W^n &= \sup_v \inf_u \left[Q(u, v) w^{n-1} + \hat{K}(u, v) \right], \\ w^n &= W^n - W_{j_0}^n \mathbf{e}. \end{aligned} \quad (3.9)$$

The condition (1.3) is no longer sufficient for convergence. For arbitrary controls $\{\hat{u}^n, \hat{v}^n\}$, let $q_{ij}^{(n)}(\hat{u}^n, \hat{v}^n; \dots; \hat{u}^1, \hat{v}^1)$ denote the i, j th element of $Q(\hat{u}^n, \hat{v}^n) \cdots Q(\hat{u}^1, \hat{v}^1)$. We now require the additional condition that there are $\epsilon > 0, j_0$, and an integer m , such that for all controls $\{\hat{u}^n, \hat{v}^n\}$ and all i ,

$$q_{ij_0}^{(m)}(\hat{u}^m, \hat{v}^m; \hat{u}^{m-1}, \hat{v}^{m-1}; \dots; \hat{u}^1, \hat{v}^1) \geq \epsilon > 0. \quad (3.10)$$

The condition is discussed below the theorem.

Theorem 3.2. w^n converges to the value $\bar{\gamma}$ of the game.

Proof. The proof is just an adaptation of that of Theorem 3.1, analogously to the way that the proof of Theorem 2.1 is an adaptation of the proof of the convergence of the classical procedure (2.2) of value iteration for the discounted cost problem. Let u^n, v^n be the selected values in (3.8) or (3.9). Then the inequalities (3.3) hold with (Q, \hat{K}) replacing (P, K) . Analogously to the development in Theorem 3.1, this and (3.10) imply (3.7) and the theorem. ■

Discussion of (3.10). Consider a one dimensional reflected diffusion on the finite interval $[A, B]$, $B > A$, and let the variance be strictly positive. Approximate this by an N -dimensional Markov chain via the methods of [9]. The reflecting states are 1 and N , which correspond to A and B , resp. If the discretization interval is small enough, then each state communicates with its immediate neighbors only, with probabilities that are bounded away from zero, uniformly in the controls. Then $\inf_{u, v, i} q_{i,2}(u, v) > 0$ and we can use any $m \geq 1$ and $j_0 = 2$ in (3.10). This is a consequence of the form of the Gauss-Seidel iteration, which connects states to those that are lower in the order of the iteration. An analogous result holds for the multidimensional case, if the diffusion being approximated is non-degenerate. See [9] for details concerning the approximation, which is the same for the game problem.

References

- [1] D.P. Bertsekas. *Dynamic Programming: Deterministic and Stochastic Models*. Prentice-Hall, Englewood Cliffs, NJ, 1987.
- [2] J. Filar and K. Vrieze. *Competitive Markov Decision Processes*. Springer-Verlag, Berlin and New York, 1996.
- [3] A. J. Hoffman and R. M. Karp. On nonterminating stochastic games. *Management Science*, 12:359–370, 1966.
- [4] H.J. Kushner. Numerical approximations for stochastic differential games: The ergodic case. to appear, *SIAM J. Control Optim.* 2003; online version available now.
- [5] H.J. Kushner. *Introduction to Stochastic Control Theory*. Holt, Rinehart and Winston, New York, 1972.
- [6] H.J. Kushner. Numerical methods for stochastic differential games. *SIAM J. Control Optim.*, pages 457–486, 2003.
- [7] H.J. Kushner and S.G. Chamberlain. Finite state stochastic games: Existence theorems and computational procedures. *IEEE Trans. on Automatic Control*, 14:248–255, 1969.
- [8] H.J. Kushner and S.G. Chamberlain. On stochastic differential games: Sufficient conditions that a given strategy be a saddle point and numerical procedures for the solution of the game. *J. Math. Anal. Appl.*, 26:560–575, 1969.
- [9] H.J. Kushner and P. Dupuis. *Numerical Methods for Stochastic Control Problems in Continuous Time*. Springer-Verlag, Berlin and New York, 1992. Second edition, 2001.
- [10] H.J. Kushner and A.J. Kleinman. Numerical methods for the solution of degenerate nonlinear equations arising in optimal stochastic control theory. *IEEE Trans. Automat. Contr.*, 13:344–353, 1968.
- [11] L. S. Shapley. Stochastic games. *Proc. Nat. Acad. Sci. U.S.A.*, 39:1095–1100, 1953.
- [12] D.J. White. Dynamic programming, Markov chains and the method of successive approximations. *J. Math. Anal Appl.*, 6:373–376, 1963.