

CRITICAL GROUPS OF SIMPLICIAL COMPLEXES

ART M. DUVAL, CAROLINE J. KLIVANS, AND JEREMY L. MARTIN

ABSTRACT. We generalize the theory of critical groups from graphs to simplicial complexes. Specifically, given a simplicial complex, we define a family of abelian groups in terms of combinatorial Laplacian operators, generalizing the construction of the critical group of a graph. We show how to realize these critical groups explicitly as cokernels of reduced Laplacians, and prove that they are finite, with orders given by weighted enumerators of simplicial spanning trees. We describe how the critical groups of a complex represent flow along its faces, and sketch another potential interpretation as analogues of Chow groups.

1. INTRODUCTION

Let G be a finite, simple, undirected, connected graph. The *critical group* of G is a finite abelian group $K(G)$ whose cardinality is the number of spanning trees of G . The critical group is an interesting graph invariant in its own right, and it also arises naturally in the theory of a discrete dynamical system with many essentially equivalent formulations — the *chip-firing game*, *dollar game*, *abelian sandpile model*, etc.— that has been discovered independently in contexts including statistical physics, arithmetic geometry, and combinatorics. There is an extensive literature on these models and their behavior: see, e.g., [3, 4, 7, 13, 20]. In all guises, the model describes a certain type of discrete flow along the edges of G . The elements of the critical group correspond to states in the flow model that are stable, but for which a small perturbation causes an instability.

The purpose of this paper is to extend the theory of the critical group from graphs to simplicial complexes. We define the higher critical groups $K_i(\Delta)$ of a finite simplicial complex Δ in terms of its simplicial chain groups, boundary maps, and combinatorial Laplacian operators. Our main result (Theorem 3.4) states that, under certain mild assumptions on the complex Δ , the groups $K_i(\Delta)$ can be expressed from a reduced version of its combinatorial Laplacian operators. It then follows from a simplicial analogue of the matrix-tree theorem [9, 10] that the orders $|K_i(\Delta)|$ of the higher critical groups are given by a torsion-weighted enumeration of higher-dimensional spanning trees (Corollary 4.2) and in terms of the eigenvalues of the Laplacian operators (Corollary 4.4). In the case that Σ is a simplicial d -sphere, we prove (Theorem 4.6) that the top-dimensional critical group $K_{d-1}(\Sigma)$ is cyclic, with order equal to the number of facets, generalizing the corresponding statement [21, 22, 23] for cycle graphs. We also give a model of

Date: January 20, 2011.

1991 Mathematics Subject Classification. Primary 05E45; Secondary 05C50, 05C21, 55U15, 57M15.

Key words and phrases. graph, simplicial complex, critical group, combinatorial Laplacian, chip-firing game, sandpile model, spanning trees.

discrete flow (Section 5) on the codimension-one faces along facets of the complex whose behavior is captured by the group structure. Finally, we outline (Section 6) an alternative interpretation of the higher critical groups as discrete analogues of the Chow groups of an algebraic variety.

The authors thank Andy Berget, Hailong Dao, Craig Huneke, Manoj Kummini, Gregg Musiker, Igor Pak, Vic Reiner, and Ken Smith for numerous helpful discussions.

2. CRITICAL GROUPS OF GRAPHS

2.1. The chip-firing game. We summarize the chip-firing game on a graph, omitting the proofs. For more details, see, e.g., Biggs [3].

Let $G = (V, E)$ be a finite, simple¹, connected, undirected graph, with $V = [n] \cup q = \{1, 2, \dots, n, q\}$ and $E = \{e_1, \dots, e_m\}$. The special vertex q is called the *bank* (or “root” or “government”). Let d_i be the degree of vertex i , i.e. the number of adjacent vertices. The chip-firing game is a discrete dynamical system whose state is described by a *configuration* vector $\mathbf{c} = (c_1, \dots, c_n) \in \mathbb{N}^n$. Each c_i is a nonnegative integer that we think of as the number of “chips” belonging to vertex i . (Note that the number c_q of chips belonging to the bank q is not part of the data of a configuration.)

Each non-root vertex is generous (it likes to donate chips to its neighbors), egalitarian (it likes all its neighbors equally), and prudent (it does not want to go into debt). Specifically, a vertex v_i is called *ready* in a configuration \mathbf{c} if $c_i \geq d_i$. If a vertex is ready, it can *fire* by giving one chip to each of its neighbors. Unlike the other vertices, the bank is a miser. As long as other vertices are firing, the bank does not fire, but just collects chips.

As more and more chips accumulate at the bank, the game eventually reaches a configuration in which no non-bank vertex can fire. Such a configuration is called *stable*. At this point, the bank finally fires, giving one chip to each of its neighbors. Unlike the other vertices, the bank is allowed to go into debt: that is, we do not require that $c_q \geq d_q$ for the bank to be able to fire.

Denote by $\mathbf{c}(x_1, \dots, x_r)$ the configuration obtained from \mathbf{c} by firing the vertices x_1, \dots, x_r in order. This sequence (which may contain repetitions) is called a *firing sequence* for \mathbf{c} if every firing is permissible: that is, for each i , either $x_i \neq q$ is ready to fire in the configuration $\mathbf{c}(x_1, \dots, x_{i-1})$, or else $x_i = q$ and $\mathbf{c}(x_1, \dots, x_{i-1})$ is stable. A configuration \mathbf{c} is called *recurrent* if there is a nontrivial firing sequence X such that $\mathbf{c}(X) = \mathbf{c}$.

A configuration is called *critical* if it is both stable and recurrent. For every starting configuration \mathbf{c} , there is a uniquely determined critical configuration $[\mathbf{c}]$ that can be reached from \mathbf{c} by some firing sequence [3, Thm. 3.8]. The *critical group* $K(G)$ is defined as the set of these critical configurations, with group law given by $[\mathbf{c}] + [\mathbf{c}'] = [\mathbf{c} + \mathbf{c}']$, where the right-hand addition is componentwise addition of vectors.

In the *abelian sandpile model* [7], grains of sand are piled at each vertex. An additional grain of sand is added to a chosen vertex, potentially causing the pile to topple. Toppling similarly distributes one grain of sand to each neighbor and is allowed precisely when a pile has at least as many grains of sand as neighbors.

¹The chip-firing game and our ensuing results can easily be extended to allow parallel edges; we assume that G is simple for the sake of ease of exposition.

In this model, if the first toppling causes other vertices to be able to topple, they do as well. One shows that the order of these subsequent fires does not matter, motivating the name “abelian”. Eventually, the system stabilizes. The sandpile model itself is the random walk on the stable configurations and the critical group is the group generated by the avalanche operator.

The critical group can also be viewed as a discrete analogue of the Picard group of an algebraic curve. This point of view goes back at least as far as the work of Lorenzini [21, 22] and was developed, using the language of divisors, by Bacher, de la Harpe, and Nagnibeda [1] (who noted that their “setting has a straightforward generalization to higher dimensional objects”). It appears in diverse combinatorial contexts including elliptic curves over finite fields (Musiker [24]), linear systems on tropical curves (Haase, Musiker and Yu [14]), and Riemann-Roch theory for graphs (Baker and Norine [2]).

2.2. The algebraic viewpoint. The critical group can alternatively be defined algebraically in terms of the Laplacian matrix.

Definition 2.1. Let G be a finite, simple, connected, undirected graph with vertices $\{1, \dots, n, q\}$. The Laplacian matrix of G is the symmetric matrix L (or, equivalently, linear self-adjoint operator) whose rows and columns are indexed by the vertices of G , with entries

$$l_{ij} = \begin{cases} d_i & \text{if } i = j, \\ -1 & \text{if } ij \in E, \\ 0 & \text{otherwise.} \end{cases}$$

Firing vertex i in the chip-firing game is equivalent to subtracting the i^{th} column of the Laplacian (ignoring the entry indexed by q) from the configuration vector \mathbf{c} . Equivalently, if $\mathbf{c}' = \mathbf{c}(x_1, \dots, x_r)$, then the configurations \mathbf{c} and \mathbf{c}' represent the same element of the cokernel of the Laplacian (that is, the quotient of \mathbb{Z}^{n+1} by the column space of L).

It is immediate from the definition of L that $L(\mathbf{1}) = \mathbf{0}$, where $\mathbf{1}$ and $\mathbf{0}$ denote the all-ones and all-zeros vectors in \mathbb{N}^{n+1} . Moreover, it is not difficult to show that $\text{rank } L = |V| - 1 = n$. In terms of homological algebra, we have a chain complex

$$(1) \quad \mathbb{Z}^{n+1} \xrightarrow{L} \mathbb{Z}^{n+1} \xrightarrow{S} \mathbb{Z} \rightarrow 0$$

where $S(\mathbf{c}) = \mathbf{c} \cdot \mathbf{1} = c_q + c_1 + \dots + c_n$. The equation $L(\mathbf{1}) = \mathbf{0}$ says that $\ker(S) \supseteq \text{im}(L)$, and the equality $\text{rank } L = n$ implies that the abelian group $\ker(S)/\text{im}(L)$ is finite (since $\ker(S) \cong \mathbb{Z}^n$).

Definition 2.2. The *critical group* of a graph G is $K(G) = \ker(S)/\text{im}(L)$.

This definition of the critical group is equivalent to that in terms of the chip-firing game [3, Thm. 4.2]. The order of the critical group is the determinant of the *reduced Laplacian* formed by removing the row and column indexed by q [3, Thm. 6.2]. By the matrix-tree theorem, this is the number of spanning trees. As we will see, the algebraic description provides a natural framework for generalizing the critical group.

3. THE CRITICAL GROUPS OF A SIMPLICIAL COMPLEX

We assume familiarity with the basic algebraic topology of simplicial complexes; see, e.g., Hatcher [16]. Let Δ be a d -dimensional simplicial complex. For $-1 \leq i \leq d$, let $C_i(\Delta; \mathbb{Z})$ be the i^{th} simplicial chain group of Δ . We denote the simplicial boundary and coboundary maps respectively by

$$\begin{aligned}\partial_{\Delta, i} &: C_i(\Delta; \mathbb{Z}) \rightarrow C_{i-1}(\Delta; \mathbb{Z}), \\ \partial_{\Delta, i}^* &: C_{i-1}(\Delta; \mathbb{Z}) \rightarrow C_i(\Delta; \mathbb{Z}),\end{aligned}$$

where we have identified cochains with chains via the natural inner product. We will abbreviate the subscripts in the notation for boundaries and coboundaries whenever no ambiguity can arise.

Let $-1 \leq i \leq d$. The i -dimensional combinatorial Laplacian² of Δ is the operator

$$L_{\Delta, i} = \partial_{i+1} \partial_{i+1}^* : C_i(\Delta; \mathbb{Z}) \rightarrow C_i(\Delta; \mathbb{Z}).$$

Combinatorial Laplacian operators seem to have first appeared in the work of Eckmann [11] on finite dimensional Hodge theory. As the name suggests, they are discrete versions of the Laplacian operators on differential forms on a Riemannian manifold. In fact, Dodziuk and Patodi [8] showed that for suitably nice triangulations of a manifold, the eigenvalues of the discrete Laplacian converge in an appropriate sense to those of the usual continuous Laplacian. For one-dimensional complexes, i.e., graphs, the combinatorial Laplacian is just the usual Laplacian matrix $L = D - A$, where D is the diagonal matrix of vertex degrees and A is the (symmetric) adjacency matrix.

In analogy to the chain complex of (1), we have the chain complex

$$C_i(\Delta; \mathbb{Z}) \xrightarrow{L} C_i(\Delta; \mathbb{Z}) \xrightarrow{\partial_i} C_{i-1}(\Delta; \mathbb{Z}),$$

where $L = L_{\Delta, i}$. (This is a chain complex because $\partial_i L = \partial_i \partial_{i+1} \partial_{i+1}^* = 0$.) We are now ready to make our main definition.

Definition 3.1. The i -dimensional critical group of Δ is

$$K_i(\Delta) := \ker \partial_i / \text{im } L = \ker \partial_i / \text{im}(\partial_{i+1} \partial_{i+1}^*).$$

Note that $K_0(\Delta)$ is just the critical group of the 1-skeleton of Δ .

3.1. Simplicial spanning trees. Our results about critical groups rely on the theory of simplicial and cellular spanning trees developed in [9], based on earlier work of Bolker [5] and Kalai [18]. Here we briefly review the definitions and basic properties, including the higher-dimensional analogues of Kirchhoff's matrix-tree theorem. For simplicity, we present the theory for simplicial complexes, the case of primary interest in combinatorics. Nevertheless, the definitions of spanning trees, their enumeration using a generalized matrix-tree theorem, and the definition and main result about critical groups are all valid in the more general setting of regular CW-complexes [10].

In order to define simplicial spanning trees, we first fix some notation concerning simplicial complexes and algebraic topology. The symbol Δ_i will denote the set of cells of dimension i . The i -dimensional skeleton $\Delta_{(i)}$ of a simplicial complex Δ is

² In other settings, our Laplacian might be referred to as the ‘‘up-down’’ Laplacian, L^{ud} , and ‘‘Laplacian’’ might mean the (total) Laplacian $L_i^{\text{tot}} = \partial_{i+1} \partial_{i+1}^* + \partial_i^* \partial_i$. We adopt the notation we do since, as in the graphical case, we only need the up-down Laplacian.

the subcomplex consisting of all cells of dimension $\leq i$. A complex is *pure* if all maximal cells have the same dimension. The i^{th} reduced homology group of Δ with coefficients in a ring R is denoted $\tilde{H}_i(\Delta, R)$. The *Betti numbers* of Δ are $\beta_i(\Delta) = \dim_{\mathbb{Q}} \tilde{H}_i(\Delta; \mathbb{Q})$. The *f-vector* is $f(\Delta) = (f_{-1}(\Delta), f_0(\Delta), \dots)$, where $f_i(\Delta)$ is the number of faces of dimension i .

Definition 3.2. Let Δ be a pure d -dimensional simplicial complex, and let $\Upsilon \subseteq \Delta$ be a subcomplex such that $\Upsilon_{(d-1)} = \Delta_{(d-1)}$. We say that Υ is a (*simplicial*) *spanning tree* of Δ if the following three conditions hold:

- (1) $\tilde{H}_d(\Upsilon, \mathbb{Z}) = 0$;
- (2) $\tilde{H}_{d-1}(\Upsilon, \mathbb{Q}) = 0$ (equivalently, $|\tilde{H}_{d-1}(\Upsilon, \mathbb{Z})| < \infty$);
- (3) $f_d(\Upsilon) = f_d(\Delta) - \beta_d(\Delta) + \beta_{d-1}(\Delta)$.

More generally, an *i-dimensional spanning tree* of Δ is a spanning tree of the i -dimensional skeleton of Δ .

In the case $d = 1$ (that is, Δ is a graph), we recover the usual definition of a spanning tree: the three conditions above say respectively that Υ is acyclic, connected, and has one more vertex than edge. Meanwhile, the 0-dimensional spanning trees of Δ are its vertices (more precisely, the subcomplexes of Δ with a single vertex), which are precisely the connected, acyclic subcomplexes of $\Delta_{(0)}$.

Just as in the graphical case, any two of the conditions of Definition 3.2 imply the third [9, Prop 3.5]. In order for Δ to have a d -dimensional spanning tree, it is necessary and sufficient that $\tilde{H}_i(\Delta; \mathbb{Q}) = 0$ for all $i < d$; such a complex is called *acyclic in positive codimension*, or APC. Note that a graph is APC if and only if it is connected.

Example 3.3. Consider the *equatorial bipyramid*, the two-dimensional simplicial complex B with vertices [5] and facets 123, 124, 125, 134, 135, 234, 235. A geometric realization of B is shown in Figure 1. A 2-SST of B can be constructed by removing two facets F, F' , provided that $F \cap F'$ contains neither of the vertices 4, 5. A simple count shows that there are 15 such pairs F, F' , so B has 15 two-dimensional spanning trees.




FIGURE 1. The equatorial bipyramid B .

A phenomenon arising only in dimension $d > 1$ is that spanning trees may have torsion: that is, $\tilde{H}_{d-1}(\Upsilon)$ can be finite but nontrivial. For example, the 2-dimensional skeleton of a 6-vertex simplex has (several) spanning trees Υ that are homeomorphic to the real projective plane, and in particular have $\tilde{H}_1(\Upsilon; \mathbb{Z}) \cong$

$\mathbb{Z}/2\mathbb{Z}$. This cannot happen in dimension 1 (i.e., for graphs), in which every spanning tree is a contractible topological space. This torsion directly affects tree enumeration in higher dimension; see Section 4.

3.2. The main theorem. Our main result gives an explicit form for the critical group $K_i(\Delta)$ in terms of a reduced Laplacian matrix. This reduced form is both more convenient for computing examples, and gives a direct connection with the simplicial and cellular generalizations of the matrix-tree theorem [9, 10]. For a general reference on the homological algebra we will need, see, e.g., Lang [19].

Let Δ be a pure, d -dimensional, APC simplicial complex, and fix $i < d$. Let Υ be an i -dimensional spanning tree of $\Delta_{(i)}$, and let $\Theta = \Delta_i \setminus \Upsilon$ (the set of i -dimensional faces of Δ *not* in Υ). Let \tilde{L} denote the reduced Laplacian obtained from L by removing the rows and columns corresponding to Υ (equivalently, by restricting L to the rows and columns corresponding to Θ).

Theorem 3.4. *Suppose that $\tilde{H}_{i-1}(\Upsilon; \mathbb{Z}) = 0$. Then*

$$K_i(\Delta) \cong \mathbb{Z}^\Theta / \text{im } \tilde{L}.$$

Proof. We will construct a commutative diagram

$$(2) \quad \begin{array}{ccccccc} 0 & \longrightarrow & \text{im } L & \longrightarrow & \ker \partial_{\Delta, i} & \longrightarrow & K_i(\Delta) \longrightarrow 0 \\ & & \downarrow f & & \downarrow g & & \downarrow h \\ 0 & \longrightarrow & \text{im } \tilde{L} & \longrightarrow & \mathbb{Z}^\Theta & \longrightarrow & \mathbb{Z}^\Theta / \text{im } \tilde{L} \longrightarrow 0 \end{array}$$

where the rows are short exact sequences with the natural inclusions and quotient maps. The map f is defined by $f(L\theta) = \tilde{L}\theta$ for all $\theta \in \Theta$ (which we will show is an isomorphism in Claim 4), and the map g is defined by $g(\theta) = \theta$ (which we will show is an isomorphism in Claim 5). In Claim 6, we will show that $f(\gamma) = g(\gamma)$ for all $\gamma \in \text{im } L$, so the left-hand square commutes. Having proven these facts, the map h is well-defined by a diagram-chase, and it is an isomorphism by the snake lemma. We organize the proof into a series of claims.

Claim 1. $\text{im } \partial_{\Delta, i} = \text{im } \partial_{\Upsilon, i}$ as \mathbb{Z} -modules.

Indeed, we have $\text{im } \partial_{\Upsilon, i} \subseteq \text{im } \partial_{\Delta, i} \subseteq \ker \partial_{\Delta, i-1} = \ker \partial_{\Upsilon, i-1}$ (the last equality because $\Upsilon_{(i-1)} = \Delta_{(i-1)}$), so there is a short exact sequence of \mathbb{Z} -modules

$$0 \rightarrow \frac{\text{im } \partial_{\Delta, i}}{\text{im } \partial_{\Upsilon, i}} \rightarrow \frac{\ker \partial_{\Upsilon, i-1}}{\text{im } \partial_{\Upsilon, i}} \rightarrow \frac{\ker \partial_{\Delta, i-1}}{\text{im } \partial_{\Delta, i}} \rightarrow 0.$$

Since Υ is a torsion-free spanning tree, the middle term $\tilde{H}_{i-1}(\Upsilon; \mathbb{Z})$ is zero. Therefore, the first term is zero as well, proving Claim 1.

Claim 2. $\text{coker } \partial_{\Upsilon, i}$ is a free \mathbb{Z} -module.

We will use some of the basic theory of projective modules [19, pp. 137–139]. The image of $\partial_{\Upsilon, i}$ is a submodule of $C_{i-1}(\Upsilon; \mathbb{Z})$, so it is free, hence projective. Therefore, the short exact sequence

$$0 \rightarrow \ker \partial_{\Upsilon, i} \rightarrow C_i(\Upsilon; \mathbb{Z}) \xrightarrow{\partial_{\Upsilon, i}} \text{im } \partial_{\Upsilon, i} \rightarrow 0$$

is split: that is, $C_i(\Upsilon; \mathbb{Z}) = \ker \partial_{\Upsilon, i} \oplus F$, where F is a free \mathbb{Z} -module. On the other hand, $\text{im } \partial_{\Upsilon, i} \subseteq \ker \partial_{\Upsilon, i-1}$, so

$$\text{coker } \partial_{\Upsilon, i} = \frac{C_{i-1}(\Upsilon; \mathbb{Z})}{\text{im } \partial_{\Upsilon, i}} = \frac{\ker \partial_{\Upsilon, i-1} \oplus F}{\text{im } \partial_{\Upsilon, i}} = \frac{\ker \partial_{\Upsilon, i-1}}{\text{im } \partial_{\Upsilon, i}} \oplus F = \tilde{H}_{i-1}(\Upsilon; \mathbb{Z}) \oplus F$$

and $\tilde{H}_{i-1}(\Upsilon; \mathbb{Z}) = 0$ by hypothesis, proving Claim 2.

Claim 3. The coboundary map $\partial_{\Upsilon, i}^*: C_{i-1}(\Upsilon; \mathbb{Z}) \rightarrow C_i(\Upsilon; \mathbb{Z})$ is surjective.

By the basic theory of finitely generated abelian groups, we may write

$$(3) \quad \partial_{\Upsilon, i} = P \left[\begin{array}{c|c} D & 0 \\ \hline 0 & 0 \end{array} \right] Q$$

where $P \in GL_{f_{i-1}}(\mathbb{Z})$, $Q \in GL_{f_i}(\mathbb{Z})$, and D is a diagonal matrix whose entries are the cyclic summands of the torsion submodule of $\text{coker } \partial_{\Upsilon, i}$. In fact, these entries are all 1 by Claim 2. Moreover, the columns of $\partial_{\Upsilon, i}$ are linearly independent over \mathbb{Q} and over \mathbb{Z} because Υ is a simplicial tree, so in fact there are no zero columns in (3). Therefore

$$\partial_{\Upsilon, i} = P \left[\begin{array}{c} I \\ 0 \end{array} \right] Q = P \left[\begin{array}{c} Q \\ 0 \end{array} \right]$$

and transposing yields

$$\partial_{\Upsilon, i}^* = [Q^T \mid 0] P^T$$

and so

$$\begin{aligned} \text{im } \partial_{\Upsilon, i}^* &= \text{im } [Q^T \mid 0] && \text{(because } P, \text{ hence } P^T, \text{ is invertible over } \mathbb{Z}) \\ &= C_i(\Upsilon; \mathbb{Z}) && \text{(because } Q, \text{ hence } Q^T, \text{ is invertible over } \mathbb{Z}). \end{aligned}$$

We have proved Claim 3.

Claim 4. $L(C_i(\Delta; \mathbb{Z})) = L(C_i(\Theta; \mathbb{Z}))$.

Choose an arbitrary chain $\gamma \in C_i(\Upsilon; \mathbb{Z})$. By Claim 3, there is a chain $\eta \in C_{i-1}(\Upsilon; \mathbb{Z})$ such that $\partial_{\Upsilon, i}^*(\eta) = \gamma$. On the other hand, $\partial_{\Delta, i}^*(\eta) = \partial_{\Upsilon, i}^*(\eta) - \theta$ for some chain $\theta \in C_i(\Theta; \mathbb{Z})$. Hence

$$L(\gamma) - L(\theta) = L(\gamma - \theta) = L(\partial_{\Upsilon, i}^*(\eta) - \theta) = L\partial_{\Delta, i}^*(\eta) = \partial_{i+1}\partial_{i+1}^*\partial_i^*(\eta) = 0$$

and so $L(\gamma) = L(\theta)$. In particular, $L(\gamma) \in L(C_i(\Theta; \mathbb{Z}))$, which proves Claim 4.

Observe that

$$L = [L(C_i(\Upsilon; \mathbb{Z})) \mid L(C_i(\Theta; \mathbb{Z}))] = \left[\begin{array}{c|c} L(C_i(\Upsilon; \mathbb{Z})) & \\ \hline & \frac{*}{\tilde{L}} \end{array} \right].$$

Thus Claim 4 says that the Θ -columns span the full column space of L . Since L is a symmetric matrix, this statement remains true if we replace ‘‘column’’ with ‘‘row’’. In particular, there is an isomorphism $f: L(C_i(\Theta; \mathbb{Z})) \rightarrow \tilde{L}(C_i(\Theta; \mathbb{Z}))$ given by deleting the Υ -rows; that is, $f(L\theta) = \tilde{L}\theta$.

By Claim 1, for each chain $\theta \in C_i(\Theta; \mathbb{Z})$, we can write

$$\partial_{\Delta, i}(\theta) = \sum_{\sigma \in \Upsilon_i} c_{\sigma\theta} \partial_{\Delta, i}(\sigma)$$

with $c_{\sigma\theta} \in \mathbb{Z}$. Therefore, the chain

$$\hat{\theta} = \theta - \sum_{\sigma \in \Upsilon_i} c_{\sigma\theta} \sigma$$

lies in $X := \ker \partial_{\Delta, i}$.

Claim 5. The set $\{\hat{\theta} : \theta \in \Theta\}$ is a \mathbb{Z} -module basis for X .

Indeed, for any $\gamma = \sum_{\sigma \in \Delta_i} a_\sigma \sigma \in X$, let

$$\gamma' = \sum_{\sigma \in \Delta_i} a_\sigma \sigma - \sum_{\sigma \in \Theta} a_\sigma \hat{\sigma} = \sum_{\sigma \in \Upsilon_i} a_\sigma \sigma + \sum_{\sigma \in \Theta} a_\sigma (\sigma - \hat{\sigma}).$$

By the previous observation, we have $\gamma' \in X \cap C_i(\Upsilon; \mathbb{Z}) = \tilde{H}_i(\Upsilon, \mathbb{Z})$. On the other hand, $\tilde{H}_i(\Upsilon, \mathbb{Z}) = 0$ (because Υ is an i -dimensional simplicial tree), so in fact $\gamma' = 0$. Therefore, $\gamma = \sum_{\sigma \in \Theta} a_\sigma \hat{\sigma}$, proving Claim 5.

Claim 6. Suppose that the action of the reduced Laplacian \tilde{L} on $C_i(\Theta; \mathbb{Z})$ is given by

$$\tilde{L}\theta = \sum_{\sigma \in \Theta} \ell_{\theta\sigma} \sigma$$

for $\theta \in \Theta$. Then $L\theta = \sum_{\sigma \in \Theta} \ell_{\theta\sigma} \hat{\sigma}$.

Indeed, the chain $L\theta - \sum_{\sigma \in \Theta} \ell_{\theta\sigma} \hat{\sigma}$ belongs both to X and to $C_i(\Upsilon; \mathbb{Z})$, so it must be zero (as in the proof of Claim 5, because Υ is an i -dimensional simplicial tree), establishing Claim 6 and completing the proof \square

Remark 3.5. Claim 1 holds for any subcomplex $\Upsilon \subseteq \Delta$ containing a torsion-free spanning tree or, more interestingly, if Υ is “torsion-minimal”, i.e., if $\tilde{H}_{i-1}(\Upsilon, \mathbb{Z}) = \tilde{H}_{i-1}(\Delta, \mathbb{Z})$. The remainder of the proof requires Υ to be torsion-free, but it should be possible to extend these methods to the case that it is torsion-minimal.

Not every APC complex need have a torsion-minimal spanning tree. For instance, for $k \in \mathbb{N}$, let $M_k = M(\mathbb{Z}/k\mathbb{Z}, 2)$ be the Moore space [16, p. 143] obtained by attaching a 2-cell to the circle S^1 by a map of degree k ; let X be the complex obtained by identifying the 1-skeletons of M_2 and M_3 (thus, X is a CW-complex with two 2-cells, one 1-cell, and one 0-cell); and let Δ be a simplicial triangulation of X (in particular, note that Δ is a regular CW-complex with the same homology as X). Then $\tilde{H}_2(\Delta; \mathbb{Z}) \cong \mathbb{Z}$ and $\tilde{H}_i(\Delta; \mathbb{Z}) = 0$ for $i < 2$. On the other hand, the simplicial spanning trees of Δ are precisely the subcomplexes Υ obtained by deleting a single facet σ (just as though Υ were a simplicial sphere, which it certainly is not—see Remark 4.5 and subsequently), and each such Υ has $\tilde{H}_1(\Upsilon; \mathbb{Z}) \cong \mathbb{Z}/3\mathbb{Z}$ or $\mathbb{Z}/2\mathbb{Z}$, according as σ is a face of M_2 or M_3 . (We thank Vic Reiner for providing this example.)

Example 3.6. We return to the bipyramid B from Example 3.3 to illustrate Theorem 3.4. We must first pick a 1-dimensional spanning tree Υ ; we take Υ to be the spanning tree with edges 12, 13, 14, 15. (In general, we must also make sure Υ is torsion-free, but this is always true for 1-dimensional trees.) Let $L = L_{B, 1} : C_1(B; \mathbb{Z}) \rightarrow C_1(B; \mathbb{Z})$ be the full Laplacian; note that L is a 9×9 matrix whose rows and columns are indexed by the edges of B . The reduced Laplacian \tilde{L}

is formed by removing the rows and columns indexed by the edges of Υ :

$$\tilde{L} = \begin{matrix} & 23 & 24 & 25 & 34 & 35 \\ \begin{matrix} 23 \\ 24 \\ 25 \\ 34 \\ 35 \end{matrix} & \begin{pmatrix} 3 & -1 & -1 & 1 & 1 \\ -1 & 2 & 0 & -1 & 0 \\ -1 & 0 & 2 & 0 & -1 \\ 1 & -1 & 0 & 2 & 0 \\ 1 & 0 & -1 & 0 & 2 \end{pmatrix} \end{matrix}.$$

The critical group $K_1(B)$ is the cokernel of this matrix, i.e., $K_1(B) \cong \mathbb{Z}^5 / \text{im } \tilde{L}$. Since \tilde{L} has full rank, it follows that $K_1(B)$ is finite; its order is $\det(\tilde{L}) = 15$.

4. THE ORDER OF THE CRITICAL GROUP

The matrix-tree theorem implies that the order of the critical group of a graph equals the number of spanning trees. In this section, we explain how this equality carries over to the higher-dimensional setting.

As before, let Δ be a pure d -dimensional simplicial complex. Let $\mathcal{T}_i(\Delta)$ denote the set of all i -dimensional spanning trees of Δ (that is, spanning trees of the i -dimensional skeleton $\Delta_{(i)}$). Define

$$\tau_i = \sum_{\Upsilon \in \mathcal{T}_i(\Delta)} |\tilde{H}_{i-1}(\Upsilon; \mathbb{Z})|^2,$$

$$\pi_i = \text{product of all nonzero eigenvalues of } L.$$

The following formulas relate the tree enumerators τ_i to the linear-algebraic invariants π_i .

Theorem 4.1 (The simplicial matrix-tree theorem). [9, Thm. 1.3] *For all $i \leq d$, we have*

$$\pi_i = \frac{\tau_i \tau_{i-1}}{|\tilde{H}_{i-2}(\Delta; \mathbb{Z})|^2}.$$

Moreover, if Υ is any spanning tree of $\Delta_{(i-1)}$, then

$$\tau_i = \frac{|\tilde{H}_{i-2}(\Delta; \mathbb{Z})|^2}{|\tilde{H}_{i-2}(\Upsilon; \mathbb{Z})|^2} \det L_\Upsilon.$$

Recall that when $d = 1$, the number $\tau_1(\Delta)$ is simply the number of spanning trees of the graph Δ , and $\tau_0(\Delta)$ is the number of vertices (i.e., 0-dimensional spanning trees). Therefore, the formulas above specialize to the classical matrix-tree theorem.

Corollary 4.2. *Let $i < d$. Suppose that $\tilde{H}_{i-1}(\Delta; \mathbb{Z}) = 0$ and that Δ has an i -dimensional spanning tree Υ such that $\tilde{H}_{i-1}(\Upsilon; \mathbb{Z}) = 0$. Then the order of the i -dimensional critical group is the torsion-weighted number of $(i+1)$ -dimensional spanning trees, i.e.,*

$$|K_i(\Delta)| = \tau_{i+1}.$$

Example 4.3. Returning again to the bipyramid B , recall that 15 is both the number of its spanning trees (Example 3.3) and the order of its 1-dimensional critical group (Example 3.6), in each case because $\det \tilde{L} = 15$.

Another formula for the orders of the critical groups of Δ is as follows. For $0 \leq j \leq d$, denote by π_j the product of the nonzero eigenvalues of the Laplacian $L_{j-1}^{ud} = \partial_j \partial_j^*$. Then Corollary 4.2, together with [10, Corollary 2.10], implies the following formula for $|K_i(\Delta)|$ as an alternating product:

Corollary 4.4. *Under the conditions of Corollary 4.2, for every $i \leq d$, we have*

$$|K_i(\Delta)| = \prod_{j=0}^i \pi_j^{(-1)^{i-j}}.$$

The condition that Δ and Υ be torsion-free is not too restrictive, in the sense that many simplicial complexes of interest in combinatorics (such as Cohen-Macaulay complexes) are torsion-free and have torsion-free spanning trees.

Remark 4.5. When every spanning tree of Δ is torsion-free, the order of the critical group is exactly the number of spanning trees. This is a strong condition on Δ , but it does hold for some complexes — notably for simplicial spheres, whose spanning trees are exactly the (contractible) subcomplexes obtained by deleting a single facet. Thus a given explicit bijection between spanning trees and elements of the critical group amounts to an abelian group structure on the set of facets of a simplicial sphere.

Determining the structure of the critical group is not easy, even for very special classes of graphs; see, e.g., [6, 17]. One of the first such results is due to Lorenzini [21, 22] and Merris [23, Example 1(1.4)], who independently noted that the critical group of the cycle graph on n vertices is \mathbb{Z}_n , the cyclic group on n elements. Simplicial spheres are the natural generalizations of cycle graphs from a tree-enumeration point of view. In fact, the theorem of Lorenzini and Merris carries over to simplicial spheres, as we now show.

Theorem 4.6. *Let Σ be a d -dimensional simplicial sphere with n facets. Then $K_{d-1}(\Sigma) \cong \mathbb{Z}_n$.*

Proof. Let $K = K_{d-1}(\Sigma)$. Remark 4.5 implies that the group K has n elements, so it is sufficient to show that it is cyclic. In what follows, we use the standard terms “facets” and “ridges” for faces of Σ of dimensions d and $d - 1$, respectively.

By definition, K is generated by $(d - 1)$ -dimensional cycles, that is, elements of $\ker \partial_{d-1}$. Since $\tilde{H}_{d-1}(\Sigma; \mathbb{Z}) = 0$, all such cycles are in fact $(d - 1)$ -dimensional boundaries of d -dimensional chains. Therefore, K is generated by the boundaries of facets, modulo the image of $L = \partial_d \partial_d^*$. We now show that for any two facets $\sigma, \sigma' \in \Sigma$, we have $\partial_d \sigma \equiv \pm \partial_d \sigma'$ modulo $\text{im } L$. This will imply that K can be generated by a single element as a \mathbb{Z} -module.

Since Σ is a sphere, it is in particular a pseudomanifold, so every ridge is in the boundary of at most two facets [25, p. 24]. Consequently, if two facets σ, σ' share a ridge ρ , then no other facet contains ρ , and we have

$$0 \equiv \partial \partial^*(\rho) = \partial(\pm \sigma \pm \sigma') = \pm(\partial \sigma \pm \partial \sigma')$$

(where \equiv means “equal modulo $\text{im } L$ ”). Hence $\partial(\sigma)$ and $\partial(\sigma')$ represent the same or opposite elements of K . Furthermore, the definition of pseudomanifold guarantees that for any two facets σ, σ' , there is a sequence of facets $\sigma = \sigma_0, \sigma_1, \dots, \sigma_k = \sigma'$ such that each σ_j and σ_{j+1} share a common ridge. Therefore, by transitivity, the boundary of any single facet generates K , as desired. \square

The condition that Σ be a simplicial sphere can be relaxed: in fact, the proof goes through for any d -dimensional pseudomanifold Σ such that $\tilde{H}_{d-1}(\Sigma; \mathbb{Z}) = 0$. On the other hand, if Σ is APC in addition to being a pseudomanifold (for example, certain lens spaces—see [16, p. 144]), then it has the rational homology type of either a sphere or a ball (because $\tilde{H}_d(\Sigma; \mathbb{Q})$ is either \mathbb{Q} or 0 ; see [25, p. 24]).

5. THE CRITICAL GROUP AS A MODEL OF DISCRETE FLOW

In this section, we describe an interpretation of the critical group in terms of flow, analogous to the chip-firing game. By definition of $K_i(\Delta)$, its elements may be represented as integer vectors $\mathbf{c} = (c_F)_{F \in \Delta_i}$, modulo an equivalence relation given by the Laplacian. These configurations are the analogues of the configurations of chips in the graph case ($i = 0$). When $i = 1$, it is natural to interpret c_F as a flow along the edge F , in the direction given by some predetermined orientation; a negative value on an edge corresponds to flow in the opposite direction. More generally, if F is an i -dimensional face, then we can interpret c_F as a generalized i -flow, again with the understanding that a negative value on a face means a i -flow in the opposite orientation. For instance, 2-flow on a triangle represents circulation around the triangle, and a negative 2-flow means to switch between clockwise and counterclockwise.

When $i = 1$, the condition $\mathbf{c} \in \ker \partial_i$ means that flow neither accumulates nor depletes at any vertex; intuitively, matter is conserved. In general, we call an i -flow *conservative* if it lies in $\ker \partial_i$. For instance, when $i = 2$, the ∂_i map converts 2-flow around a single triangle into 1-flow along the three edges of its boundary in the natural way; for a 2-flow on Δ to be conservative, the sum of the resulting 1-flows on each edge must cancel out, leaving no net flow along any edge. In general, the sum of (the boundaries of) all the i -dimensional flows surrounding an $(i-1)$ -dimensional face must cancel out along that face.

That the group $K_i(\Delta)$ is a quotient by the image of the Laplacian means that two configurations are equivalent if they differ by an integer linear combination of Laplacians applied to i -dimensional faces. This is analogous to the chip-firing game, where configurations are equivalent when it is possible to get from one to the other by a series of chip-firings, each of which corresponds to adding a column vector of the Laplacian. When $i = 1$, it is easy to see that firing an edge e (adding the image of its Laplacian to a configuration) corresponds to diverting one unit of flow around each triangle containing e (see Example 5.1). More generally, to fire an i -face F means to divert one unit of i -flow from F around each $(i+1)$ -face containing F .

By Theorem 3.4, we may compute the critical group as \mathbb{Z}^Θ modulo the image of the reduced Laplacian. In principle, passing to the reduced Laplacian means ignoring the i -flow along each facet of an i -dimensional spanning tree Υ . In the graph case ($i = 0$), this spanning tree is simply the bank vertex. The higher-dimensional generalization of this statement is that the equivalence class of a configuration \mathbf{c} is determined by the subvector $(c_F)_{F \in \Delta \setminus \Upsilon}$.

A remaining open problem is to identify the higher-dimensional “critical configurations”, i.e., a set of stable and recurrent configurations that form a set of coset representatives for the critical group. Recall that in the chip-firing game, when vertex i fires, every vertex other than i either gains a chip or stays unchanged. Therefore, we can define stability simply by the condition $\mathbf{c}_i < \deg(i)$ for every non-bank vertex i . On the other hand, when a higher-dimensional face fires, the flow along nearby faces can actually decrease. Therefore, it is not as easy to define stability. For instance, one could try to define stability by the condition that no face can fire without forcing some face (either itself or one of its neighbors) into debt. However, with this definition, there are some examples (such as the 2-skeleton of the tetrahedron) for which some of the cosets of the Laplacian admit more than one critical configuration. Therefore, it is not clear how to choose a canonical

set of coset representatives analogous to the critical configurations of the graphic chip-firing game.

Example 5.1. We return once again to the bipyramid B , and its 1-dimensional spanning tree Υ with edges 12, 13, 14, 15. If we pick 1-flows on $\Theta = \Delta_1 \setminus \Upsilon$ as shown in Figure 2(a), it is easy to compute that we need 1-flows on Υ as shown in Figure 2(b) to make the overall flow 1-conservative. Since Theorem 3.4 implies we can always pick flows on Υ to make the overall flow 1-conservative, we only show flows on Θ in subsequent diagrams.

If we fire edge 23, we get the configuration shown in Figure 2(c). One unit of flow on edge 23 has been diverted across face 234 to edges 24 and 34, and another unit of flow has been diverted across face 235 to edges 25 and 35. Note that the absolute value of flow on edge 25 has actually decreased, because of its orientation relative to edge 23. If we subsequently fire edge 24, we get the configuration shown in Figure 2(d). One unit of flow on edge 24 has been diverted across face 234 to edges 23 and 34, and another unit of flow has been diverted across face 124 to edges 12 and 14 (and out of the diagram of Θ).




FIGURE 2. Conservative 1-flows and firings

6. CRITICAL GROUPS AS CHOW GROUPS

An area for further research is to interpret the higher-dimensional critical groups of a simplicial complex Δ as simplicial analogues of the Chow groups of an algebraic variety. (For the algebraic geometry background, see, e.g., [15, Appendix A] or [12].) We regard Δ as the discrete analogue of a d -dimensional variety, so that divisors correspond to formal sums of codimension-1 faces. Even more generally, algebraic cycles of dimension i correspond to simplicial i -chains. The critical group $K_i(\Delta)$ consists of closed i -chains modulo conservative flows (in the language of Section 5) is thus analogous to the Chow group of algebraic cycles modulo rational equivalence. This point of view has proved fruitful in the case of graphs [1, 2, 14, 21, 22]. In order to develop this analogy fully, the next step is to define a ring structure on $\bigoplus_{i \geq 0} K_i(\Delta)$ with a ring structure analogous to that of the Chow ring. The goal is to define a “critical ring” whose multiplication encodes a simplicial version of intersection theory on Δ .

REFERENCES

- [1] Roland Bacher, Pierre de la Harpe, and Tatiana Nagnibeda, The lattice of integral flows and the lattice of integral cuts on a finite graph, *Bull. Soc. Math. France* **125** (1997), no. 2, 167–198.
- [2] Matthew Baker and Serguei Norine, Riemann-Roch and Abel-Jacobi theory on a finite graph, *Adv. Math.* **215** (2007), no. 2, 766–788.

- [3] Norman Biggs, Chip-firing and the critical group of a graph, *J. Alg. Combin.* **9** (1999), 25–45.
- [4] Anders Björner, László Lovász, and Peter Shor, Chip-firing games on graphs, *European J. Combin.* **12** (1991), no. 4, 283–291.
- [5] Ethan Bolker, Simplicial geometry and transportation polytopes, *Trans. Amer. Math. Soc.* **217** (1976), 121–142.
- [6] Hans Christianson and Victor Reiner, The critical group of a threshold graph. *Linear Algebra Appl.* **349** (2002), 233–244.
- [7] Deepak Dhar, Self-organized critical state of sandpile automaton models, *Phys. Rev. Lett.* **64** (1990), no. 14, 1613–1616.
- [8] Jozef Dodziuk and Vijay Kumar Patodi, Riemannian structures and triangulations of manifolds, *J. Indian Math. Soc. (N.S.)* **40** (1976), no. 1–4, 1–52 (1977).
- [9] Art M. Duval, Caroline J. Klivans, and Jeremy L. Martin, Simplicial matrix-tree theorems, *Trans. Amer. Math. Soc.* **361** (2009), no. 11, 6073–6114.
- [10] Art M. Duval, Caroline J. Klivans, and Jeremy L. Martin, Cellular spanning trees and Laplacians of cubical complexes, preprint, arXiv:0908.1956, to appear in *Adv. Appl. Math.*
- [11] Beno Eckmann, Harmonische Funktionen und Randwertaufgaben in einem Komplex, *Comment. Math. Helv.* **17** (1945), 240–255.
- [12] William Fulton, *Intersection Theory*, 2nd ed., Springer-Verlag, Berlin, 1998.
- [13] Chris Godsil and Gordon Royle, *Algebraic Graph Theory*, Graduate Texts in Mathematics 207, Springer-Verlag, New York, 2001.
- [14] Christian Haase, Gregg Musiker, and Josephine Yu, Linear systems on tropical curves, preprint, arXiv:0909.3685.
- [15] Robin Hartshorne, *Algebraic geometry*, Graduate Texts in Mathematics, 52, Springer-Verlag, New York-Heidelberg, 1977.
- [16] Allen Hatcher, *Algebraic Topology*. Cambridge University Press, Cambridge, 2002; available online at <http://www.math.cornell.edu/~hatcher/AT/ATpage.html>
- [17] Brian Jacobson, Andrew Niedermaier, and Victor Reiner, Critical groups for complete multipartite graphs and Cartesian products of complete graphs, *J. Graph Theory* **44** (2003), no. 3, 231–250.
- [18] Gil Kalai, Enumeration of \mathbf{Q} -acyclic simplicial complexes, *Israel J. Math.* **45**, no. 4 (1983), 337–351.
- [19] Serge Lang, *Algebra*, 3rd ed., Addison-Wesley, 1993.
- [20] Lionel Levine and James Propp, What is . . . a sandpile?, *Notices Amer. Math. Soc.* **57**, no. 8, 976–979.
- [21] Dino J. Lorenzini, Arithmetical graphs, *Math. Ann.* **285**, no. 3 (1989), 481–501.
- [22] Dino J. Lorenzini, A finite group attached to the Laplacian of a graph, *Discrete Math.* **91**, no. 3 (1991), 277–282.
- [23] Russell Merris, Unimodular equivalence of graphs, *Lin. Alg. Appl.* **173** (1992), 181–189.
- [24] Gregg Musiker, The critical groups of a family of graphs and elliptic curves over finite fields, *J. Algebraic Combin.* **30** (2009), no. 2, 255–276.
- [25] Richard P. Stanley, *Combinatorics and Commutative Algebra*, 2nd ed., Birkhäuser, 1996.

DEPARTMENT OF MATHEMATICAL SCIENCES, UNIVERSITY OF TEXAS AT EL PASO

DEPARTMENTS OF MATHEMATICS AND COMPUTER SCIENCE, THE UNIVERSITY OF CHICAGO

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF KANSAS