Dissipative Particle Dynamics: Foundation, Evolution and Applications Lecture 2: Theoretical foundation and parameterization







George Em Karniadakis

Division of Applied Mathematics, Brown University & Department of Mechanical Engineering, MIT & Pacific Northwest National Laboratory, CM4

The CRUNCH group: www.cfm.brown.edu/crunch



Outline

- 1. Background
- 2. Fluctuation-dissipation theorem
- 3. Parameterization
 - Static properties
 - Dynamic properties (kinetic theory)
- 4. DPD ----> Navier-Stokes
- 5. Navier-Stokes ----> (S)DPD
- 6. Microscopic ----> DPD
 - Mori-Zwanzig formalism



Outline

1. Background

- 2. Fluctuation-dissipation theorem
- 3. Parameterization
 - Static properties
 - Dynamic properties (kinetic theory)
- 4. DPD ----> Navier-Stokes
- 5. Navier-Stokes ----> (S)DPD
- 6. Microscopic ----> DPD
 - Mori-Zwanzig formalism



1. Background

- Molecular dynamics (e.g. Lennard-Jones):
 - Lagrangian nature
 - Stiff force
 - Atomic time step

(Allen&Tildesley, Oxford Uni. Press, 1989)



- Coarse-grained (1980s): Lattice gas automata
 - Mesoscopic collision rules
 - Grid based particles

(Frisch et al, PRL, 1986)





Mesoscale + Langrangian?

 Physics intuition: Let particles represent clusters of molecules and interact via pair-wise forces

$$\vec{\mathbf{F}}_{i} = \sum_{j \neq i} \left(\vec{\mathbf{F}}_{ij}^{C} + \vec{\mathbf{F}}_{ij}^{R} / \sqrt{dt} + \vec{\mathbf{F}}_{ij}^{D} \right)$$

Conditions:

- Conservative force is softer than Lennard-Jones
- System is thermostated by two forces $\vec{\mathbf{F}}^{R}$, $\vec{\mathbf{F}}^{D}$
- Equation of motion is Lagrangian as:

$$d\vec{\mathbf{r}}_{i} = \vec{\mathbf{v}}_{i}dt \quad d\vec{\mathbf{v}}_{i} = \vec{F}_{i}dt$$

This innovation is named as DPD method!



Hoogerbrugge & Koelman, EPL, 1992

Outline

1. Background

- 2. Fluctuation-dissipation theorem
- 3. Parameterization
 - Static properties
 - Dynamic properties (kinetic theory)
- 4. DPD ----> Navier-Stokes
- 5. Navier-Stokes ----> (S)DPD
- 6. Microscopic ----> DPD
 - Mori-Zwanzig formalism



2. Fluctuation-dissipation theorem

•Langevin equations (SDEs)

$$\begin{cases} \mathrm{d}\boldsymbol{r}_{i} = \frac{\boldsymbol{p}_{i}}{m_{i}} \mathrm{d}t \\ \mathrm{d}\boldsymbol{p}_{i} = \left[\sum_{j \neq i} \boldsymbol{F}_{ij}^{\mathrm{C}}(\boldsymbol{r}_{ij}) + \sum_{j \neq i} -\gamma \omega_{\mathrm{D}}(\boldsymbol{r}_{ij})(\boldsymbol{e}_{ij} \cdot \boldsymbol{v}_{ij}) \boldsymbol{e}_{ij}\right] \mathrm{d}t + \sum_{j \neq i} \sigma \omega_{\mathrm{R}}(\boldsymbol{r}_{ij}) \boldsymbol{e}_{ij} \mathrm{d}W_{ij} \end{cases}$$

With $dW_{ij} = dW_{ji}$ the independent Wiener increment: $dW_{ij}dW_{i'j'} = (\delta_{ii'}\delta_{jj'} + \delta_{ij'}\delta_{ji'}) dt$ • Corresponding Fokker-Planck equation (FPE) $\partial_t \rho(r, p; t) = L_{C}\rho(r, p; t) + L_{D}\rho(r, p; t)$

$$\begin{split} L_{\mathrm{C}}\rho(r,\,p;\,t) &\equiv -\left[\sum_{i}\frac{\boldsymbol{p}_{i}}{m}\frac{\partial}{\partial\boldsymbol{r}_{i}} + \sum_{i,\,j\neq\,i}\boldsymbol{F}_{ij}^{\mathrm{C}}\frac{\partial}{\partial\boldsymbol{p}_{i}}\right]\rho(r,\,p;\,t) \\ L_{\mathrm{D}}\rho(r,\,p;\,t) &\equiv \sum_{i,\,j\neq\,i}\boldsymbol{e}_{ij}\frac{\partial}{\partial\boldsymbol{p}_{i}}\left[\gamma\omega_{\mathrm{D}}(r_{ij})(\boldsymbol{e}_{ij}\cdot\boldsymbol{v}_{ij}) + \frac{\sigma^{2}}{2}\omega_{\mathrm{R}}^{2}(r_{ij})\boldsymbol{e}_{ij}\left(\frac{\partial}{\partial\boldsymbol{p}_{i}} - \frac{\partial}{\partial\boldsymbol{p}_{j}}\right)\right]\rho(r,\,p;\,t) \end{split}$$

2. Fluctuation-dissipation theorem

•Gibbs distribution: steady state solution of FPE

$$\varphi^{\text{eq}}(r, p) = \frac{1}{Z} \exp\left[-H(r, p)/k_{\text{B}}T\right] = \frac{1}{Z} \exp\left[-\left(\sum_{i} \frac{p_{i}^{2}}{2m_{i}} + V(r)\right)/k_{\text{B}}T\right]$$

Conservative Potential $\mathbf{F}^{C} = -\nabla \mathbf{V}(\mathbf{r})$ $L_{C}\rho^{eq} = 0$

Require $L_D \rho^{eq} = 0$ Energy dissipation and generation balance

DPD version of fluctuation-dissipation theorem $\omega_{\rm R}(r) = \omega_{\rm D}^{1/2}(r) \qquad \sigma = (2k_{\rm B}T\gamma)^{1/2}$

DPD can be viewed as canonical ensemble (NVT) Espanol, EPL, 1995



Outline

- 1. Background
- 2. Fluctuation-dissipation theorem
- 3. Parameterization
 - Static properties
 - Dynamic properties (kinetic theory)
- 4. DPD ----> Navier-Stokes
- 5. Navier-Stokes ----> (S)DPD
- 6. Microscopic ----> DPD
 - Mori-Zwanzig formalism



3. Parameterization

(How to choose simulation parameters?)

Strategy: match DPD thermodynamics to atomistic system

I. How to choose repulsion parameter?

Match the static thermo-properties, i.e.,

Isothermal compressibility (water)

Mixing free energy, Surface tension (polymer blends)

II. How to choose dissipation (or fluctuation) parameter?

Match the dynamic thermo-properties, i.e.,

Self-diffusion coefficient, kinematic viscosity (however, can not match both easily) Schmidt number $Sc = \nu/D$ usually lower than atomic fluid



Repulsion parameter for water?

Equation of state: self and pair contributions

$$p = \rho k_{\mathrm{B}} T + \frac{1}{3V} \left\langle \sum_{j > i} \left(\mathbf{r}_{i} - \mathbf{r}_{j} \right) \cdot \mathbf{f}_{i} \right\rangle = \rho k_{\mathrm{B}} T + \frac{1}{3V} \left\langle \sum_{j > i} \left(\mathbf{r}_{i} - \mathbf{r}_{j} \right) \cdot \mathbf{F}_{ij}^{\mathrm{C}} \right\rangle = \rho k_{\mathrm{B}} T + \frac{2\pi}{3} \rho^{2} \int_{0}^{1} rf(r)g(r)r^{2} dr$$



Repulsion parameter for polymers



Friction parameters for simple fluids

Simple argument by Groot&Warren, JChemPhys., 1997

Consider an uniform linear flow $v_{\alpha} = e_{\alpha\beta}r_{\beta}$

Dissipative contribution to stress

$$\sigma_{\alpha\beta} = \frac{1}{V} \left\langle \sum_{i>j} r_{ij\alpha} \mathbf{F}_{ij\beta}^{\mathsf{D}} \right\rangle = \frac{\rho^2}{2} \int d^3 \mathbf{r} \ \gamma w^{\mathsf{D}}(r) r_{\alpha} \hat{r}_{\beta} \hat{r}_{\gamma} r_{\delta} \ e_{\gamma\delta} = \frac{2 \pi \gamma \rho^2}{15} \int_0^\infty dr \ r^4 w^{\mathsf{D}}(r) [e_{\alpha\beta} + e_{\beta\alpha} + \delta_{\alpha\beta} e_{\gamma\gamma}]$$

Dissipative viscosity $\eta^{\mathsf{D}} = \frac{2 \pi \gamma \rho^2}{15} \int_0^\infty dr \ r^4 w^{\mathsf{D}}(r)$

ignore conservative forces, average out other particle velocities

$$\begin{aligned} \frac{d\mathbf{v}_{i}}{dt} + \frac{\mathbf{v}_{i}}{\tau} &= \mathbf{F}^{\mathbb{R}} \\ \frac{1}{\tau} &= \sum_{j \neq i} \gamma w^{\mathbb{D}}(r_{ij}) \frac{\hat{\mathbf{r}}_{ij} \cdot \hat{\mathbf{r}}_{ij}}{3} &= \frac{4\pi\gamma\rho}{3} \int_{0}^{\infty} dr \ r^{2}w^{\mathbb{D}}(r) \\ \mathbf{F}^{\mathbb{R}} \rangle &= 0, \quad \langle \mathbf{F}^{\mathbb{R}}(t) \cdot \mathbf{F}^{\mathbb{R}}(t') \rangle = 4\pi\sigma^{2}\rho \int_{0}^{\infty} dr \ r^{2}[w^{\mathbb{R}}(r)]^{2} \,\delta(t-t') \end{aligned}$$

Kinetic theory: dynamic properties

Marsh et al, EPL&PRE, 1997

Single-particle and pair distribution functions

 $f(x,t) = \langle \sum_i \delta(x-x_i(t)) \rangle \qquad f^{(2)}(x,x',t) = \langle \sum_{i \neq j} \delta(x-x_i(t)) \delta(x'-x_j(t)) \rangle$

assume molecular chaos, i.e. $f^{(2)}(x,x',t) = f(x,t)f(x',t)$

Fokker-Planck-Boltzmann equation

$$\partial_t f(x) + \boldsymbol{v} \cdot \boldsymbol{\nabla} f(x) = I(f)$$

with collision ter $I(f) = \partial \cdot \int \mathrm{d}x' \gamma(x,x') f(x') f(x) + \frac{1}{2} \partial \partial$: $\int \mathrm{d}x' \sigma(x,x') \sigma(x,x') f(x') f(x) = \partial \cdot \int \mathrm{d}x' \gamma(x,x') f(x') f(x) + \frac{1}{2} \partial \partial \tau$

Integration of FPB over v yields continuity equation $\partial_t n = -\nabla \cdot nu$.

Multiplying FPB by v and integrate over v yields momentum equation

$$\partial_t \rho \boldsymbol{u} = -\boldsymbol{\nabla} \cdot \int \mathrm{d} \boldsymbol{v} \boldsymbol{v} \boldsymbol{v} f(x) - m \int \mathrm{d} \boldsymbol{v} \mathrm{d} x' \boldsymbol{\gamma}(x, x') f(x') f(x) \equiv -\boldsymbol{\nabla} \cdot (\rho \boldsymbol{u} \boldsymbol{u} + \boldsymbol{\Pi}_\mathrm{K} + \boldsymbol{\Pi}_\mathrm{D})$$

 $\begin{array}{ll} \eta_{\rm D}=mn\omega_0\langle R^2\rangle_w/2(d+2), & \zeta_{\rm D}=mn\omega_0\langle R^2\rangle_w/2d\\ \text{Compare with NS}\\ \text{equation} & \eta_{\rm K}=n\theta_0/2\omega_0; & \zeta_{\rm K}=n\theta_0/d\omega_0; \end{array}$



Outline

- 1. Background
- 2. Fluctuation-dissipation theorem
- 3. Parameterization
 - Static properties
 - Dynamic properties (kinetic theory)
- 4. DPD ----> Navier-Stokes
- 5. Navier-Stokes ----> (S)DPD
- 6. Microscopic ----> DPD
 - Mori-Zwanzig formalism





Espanol, PRE, 1995

Stochastic differential equations

•DPD equations of motion

$$d\mathbf{r}_i = rac{\mathbf{p}_i}{m_i} dt,$$

$$d\mathbf{p}_{i} = \left[\sum_{j \neq i} \mathbf{F}_{ij}^{C}(\mathbf{r}_{ij}) + \sum_{j \neq i} -\gamma \omega(r_{ij})(\mathbf{e}_{ij} \cdot \mathbf{v}_{ij})\mathbf{e}_{ij}\right] dt$$

$$+\sum_{j\neq i}\sigma\omega^{1/2}(r_{ij})\mathbf{e}_{ij}dW_{ij},$$



Fokker-Planck equation

Evolution of probability density in phase space

Conservative/Liouville operatorDissipative and random operators

$$\partial_t \rho(r, p; t) = L_{\rm C} \rho(r, p; t) + L_{\rm D} \rho(r, p; t)$$

$$\begin{cases} L_{\mathrm{C}}\rho(r,\,p;\,t) \equiv -\left[\sum_{i}\frac{\boldsymbol{p}_{i}}{m}\frac{\partial}{\partial\boldsymbol{r}_{i}} + \sum_{i,j\neq i}\boldsymbol{F}_{ij}^{\mathrm{C}}\frac{\partial}{\partial\boldsymbol{p}_{i}}\right]\rho(r,\,p;\,t) \\ L_{\mathrm{D}}\rho(r,\,p;\,t) \equiv \sum_{i,j\neq i}\boldsymbol{e}_{ij}\frac{\partial}{\partial\boldsymbol{p}_{i}}\left[\gamma\omega_{\mathrm{D}}(r_{ij})(\boldsymbol{e}_{ij}\cdot\boldsymbol{v}_{ij}) + \frac{\sigma^{2}}{2}\omega_{\mathrm{R}}^{2}(r_{ij})\boldsymbol{e}_{ij}\left(\frac{\partial}{\partial\boldsymbol{p}_{i}} - \frac{\partial}{\partial\boldsymbol{p}_{j}}\right)\right]\rho(r,\,p;\,t) \end{cases}$$



Mori projection (linearized hydrodynamics)

• Relevant hydrodynamic variables to keep

$$\begin{split} \delta \rho_{\mathbf{r}} &= \sum_{i} m \delta(\mathbf{r} - \mathbf{r}_{i}) - \rho_{0}, \\ \mathbf{g}_{\mathbf{r}} &= \sum_{i} \mathbf{p}_{i} \delta(\mathbf{r} - \mathbf{r}_{i}), \\ \delta e_{\mathbf{r}} &= \sum_{i} \left[\frac{p_{i}^{2}}{2m} + \frac{1}{2} \sum_{j \neq i} \phi_{ij} \right] \delta(\mathbf{r} - \mathbf{r}_{i}) - e_{0}, \end{split}$$

•Equilibrium averages vanish



Mori projection

Navier-Stokes

$$\partial_t \mathbf{g}(\mathbf{r},t) = -c_0^2 \nabla \delta \rho(\mathbf{r},t) + \eta \nabla^2 \mathbf{v}(\mathbf{r},t)$$

$$+\left(\zeta-rac{2\eta}{3}
ight)oldsymbol{
abla}[oldsymbol{
abla}\cdot\mathbf{v}(\mathbf{r},t)]$$

Sound speed

$$c_0^2 = \left. rac{\partial p}{\partial
ho}
ight|_T$$



Espanol, PRE, 1995

Mori projection

• Stress tensor via Irving-Kirkwood formula:

$$\Sigma^{C} = \int d^{3}\mathbf{r}\sigma_{\mathbf{r}}^{C} = \sum_{i} \frac{\mathbf{p}_{i}}{m}\mathbf{p}_{i} + \sum_{ij} (\mathbf{r}_{i} - \mathbf{r}_{j})\mathbf{F}_{ij}^{C},$$

$$\Sigma^{D} = \int d^{3}\mathbf{r}\sigma_{\mathbf{r}}^{D} = \sum_{ij} (\mathbf{r}_{i} - \mathbf{r}_{j})\mathbf{F}_{ij}^{D}$$

$$= -\gamma \sum_{ij} (\mathbf{r}_i - \mathbf{r}_j) \omega_{ij} (\mathbf{e}_{ij} \cdot \mathbf{v}_{ij}) \mathbf{e}_{ij}.$$

- Contributions:
 - Conservative force
 - Dissipative force



Mori projection

Viscosities via with Green-Kubo formulas

> Shear viscosity η and bulk viscosity ζ

$$\begin{split} \eta^C &= \beta \int_0^\infty du \frac{1}{V} [\Sigma^C_{\mu\nu}(u), \mathcal{Q}\Sigma^C_{\mu\nu}], \\ \left(\zeta^C - \frac{2}{3}\eta^C\right) &= \beta \int_0^\infty du \frac{1}{V} [\Sigma^C_{\mu\mu}(u), \mathcal{Q}\Sigma^C_{\nu\nu}], \\ \eta^D &= \beta \int_0^\infty du \frac{1}{V} [\Sigma^D_{\mu\nu}(u), \mathcal{Q}\Sigma^D_{\mu\nu}], \\ \left(\zeta^D - \frac{2}{3}\eta^D\right) &= \beta \int_0^\infty du \frac{1}{V} [\Sigma^D_{\mu\mu}(u), \mathcal{Q}\Sigma^D_{\nu\nu}], \end{split}$$

 \bullet Note the squared dependence of viscosity on γ



Outline

- 1. Background
- 2. Fluctuation-dissipation theorem
- 3. Parameterization
 - Static properties
 - Dynamic properties (kinetic theory)
- 4. DPD ----> Navier-Stokes
- 5. Navier-Stokes ----> (S)DPD
- 6. Microscopic ----> DPD
 - Mori-Zwanzig formalism



5. Navier-Stokes ---> (S)DPD

Story begins with

smoothed particle hydrodynamics (SPH) method

•Originally invented for Astrophysics (Lucy. 1977, Gingold&Monaghan, 1977)

•Popular since 1990s for physics on earth (Monaghan, 2005)



SPH 1st step: kernel approximation

 $A(\mathbf{r})$: function of spatial coordinates

integral interpolant:

$$A_{I}(\mathbf{r}) = \int A(\mathbf{r}') W(\mathbf{r} - \mathbf{r}', h) d\mathbf{r}',$$

where weighting function/kernel W: (Monaghan, RepProgPhys 2005)

$$\lim_{h\to 0} W(\mathbf{r}-\mathbf{r}',h) = \delta(\mathbf{r}-\mathbf{r}'), \quad \int W(\mathbf{r}-\mathbf{r}',h)d\mathbf{r}' = 1$$

Gaussian; cubic spline; quintic spline ... (Morris et al, JComputPhys 1997)
 h > 0: kernel error

$$A(\mathbf{r}) = A_I(\mathbf{r}) + E_1(h)$$



SPH 2nd step: particle approximation

• summation form $(r_c = 3h)$:

$$A_{S}(\mathbf{r}) = \sum_{j} \frac{A_{j}}{d_{j}} W(\mathbf{r} - \mathbf{r}_{j}, h)$$
$$\nabla A_{S}(\mathbf{r}) = \sum_{j} \frac{A_{j}}{d_{j}} \nabla W(\mathbf{r} - \mathbf{r}_{j}, h)$$
$$\dots = \dots$$



compact support: cell list

(Español&Revenga, PRE 2003)

• $\Delta x > 0$: summation error $A_I(\mathbf{r}) = A_S(\mathbf{r}) + \frac{E_2(\Delta x/h)}{E_2(\Delta x/h)}$

• $A(\mathbf{r}) = A_S(\mathbf{r}) + E_1(h) + E_2(\Delta x/h)$ (Quinlan et al., IntJNumerMethEng 2006)

 Error estimated for particles on grid
 Actual error depends on configuration of particles (Price, JComputPhys. 2012)



SPH: isothermal Navier-Stokes

Continuity equation

$$d_i = \frac{\rho_i}{m_i} = \sum_j W_{ij}, \quad \dot{\mathbf{r}}_i = \mathbf{v}_i$$

Momentum equation

$$\begin{split} m_i \dot{\mathbf{v}}_i &= -\sum_{j \neq i} \left(\frac{\bar{p}_{ij}}{d_i^2} + \frac{\bar{p}_{ij}}{d_j^2} \right) \frac{\partial W}{\partial r_{ij}} \mathbf{e}_{ij} + \sum_{j \neq i} \eta \left(\frac{1}{d_i^2} + \frac{1}{d_j^2} \right) \frac{\partial W}{\partial r_{ij}} \frac{\mathbf{v}_{ij}}{r_{ij}} + \mathbf{F}_i^{Ext} \\ &= \sum_{j \neq i} \left(\mathbf{F}_{ij}^C + \mathbf{F}_{ij}^D \right) + \mathbf{F}_i^{Ext} \end{split}$$

•Input equation of state: pressure and density Hu&Adams, JComputPhys. 2006



SPH: add Brownian motion

• Momentum with fluctuation (Espanol&Revenga, 2003)

$$m_i \dot{\mathbf{v}}_i = \sum_{j \neq i} \left(\mathbf{F}_{ij}^C + \mathbf{F}_{ij}^D + \mathbf{F}_{ij}^R / \sqrt{dt} \right) + \mathbf{F}_i^{Ext}$$

•Cast dissipative force in GENERIC \rightarrow random force

$$\mathbf{F}_{ij}^{R} = \left[\frac{-4k_{B}T\eta}{r_{ij}}\left(\frac{1}{d_{i}^{2}} + \frac{1}{d_{j}^{2}}\right)\frac{\partial W}{\partial r_{ij}}\right]^{1/2}d\overline{\mathbf{W}}_{ij} \cdot \mathbf{e}_{ij}$$
$$d\overline{\mathbf{W}}_{ij} = \left(d\mathbf{W}_{ij} + d\mathbf{W}_{ij}^{T}\right)/2 - tr[d\mathbf{W}_{ij}]\mathbf{I}/D$$

•dW is an independent increment of Wiener process Espanol&Revenga, PRE, 2003

SPH + fluctuations = SDPD

•Discretization of Landau-Lifshitz's fluctuating hydrodynamics (Landau&Lifshitz, 1959)

•Fluctuation-dissipation balance on discrete level

Same numerical structure as original DPD formulation

$$m_i \dot{\mathbf{v}}_i = \sum_{j \neq i} \left(\mathbf{F}_{ij}^C + \mathbf{F}_{ij}^D + \mathbf{F}_{ij}^R / \sqrt{dt} \right) + \mathbf{F}_i^{Ext}$$



GENERIC framework (part 1)

(General equation for nonequilibrium reversible-irreversible coupling)

•Dynamic equations of a deterministic system:

$$\frac{dx}{dt} = L \frac{\delta E}{\delta x} + M \frac{\delta S}{\delta x}$$
State variables x: position, velocity, energy/entropy
E(x): energy/ S(x): entropy
L and M are linear operators/matrices and
represent reversible and irreversible dynamics

• First and second Laws of thermodynamics

$$L\frac{\delta S}{\delta x} = 0$$
 $M\frac{\delta E}{\delta x} = 0$

•For any dynamic invariant variable I, e.g, linear momentum

if
$$\frac{\partial I}{\partial x}L\frac{\partial E}{\partial x}=0$$
, $\frac{\partial I}{\partial x}M\frac{\partial S}{\partial x}=0$, then $\dot{I}=0$

Grmela&Oettinger, PRE, 1997; Oettinger&Grmela, PRE, 1997



GENERIC framework (part 2)

(General equation for nonequilibrium reversible-irreversible coupling) • Dynamic equations of a stochastic system:

$$dx = \left[L \frac{\partial E}{\partial x} + M \frac{\partial S}{\partial x} + k_B \frac{\partial}{\partial x} M \right] dt + d\tilde{x}$$
Last term is thermal fluctuations

•Fluctuation-dissipation theorem: compact form

 $d\tilde{x}d\tilde{x}^T = 2k_BMdt$

No Fokker-Planck equation needs to be derived

Model construction becomes simple linear algebra

Grmela&Oettinger, PRE, 1997; Oettinger&Grmela, PRE, 1997



Outline

- 1. Background
- 2. Fluctuation-dissipation theorem
- 3. Parameterization
 - Static properties
 - Dynamic properties (kinetic theory)
- 4. DPD ----> Navier-Stokes
- 5. Navier-Stokes ----> (S)DPD
- 6. Microscopic ----> DPD
 - Mori-Zwanzig formalism



Coarse-graining: Voronoi tessellation

•Procedure:

- 1. Partition of particles of molecular dynamics
- 2. Measuring fluxes at edges
- 3. Update center of mass
- 4. Repeat 1, 2 and 3
- 5. Ensemble average interacting forces between neighboring Voronoi cells: similarly as DPD pairwise interactions

$$\frac{dM_k}{dt} = \sum_l \dot{M}_{kl} \equiv \sum_i f_{kl}(\mathbf{x}_i) m(\mathbf{v}'_i \cdot \mathbf{r}_{kl} + \mathbf{x}'_i \cdot \mathbf{U}_{kl}),$$

$$\frac{d\mathbf{P}_k}{dt} = M_k \mathbf{g} + \sum_l \dot{M}_{kl} \frac{\mathbf{U}_k + \mathbf{U}_l}{2} + \sum_{li} f_{kl}(\mathbf{x}_i) \mathbf{\Pi}'_i \cdot \mathbf{r}_{kl},$$

$$\frac{dE_k}{dt} = \sum_l \frac{\dot{M}_{kl}}{2} \left(\frac{\mathbf{U}_{kl}}{2}\right)^2 + \sum_{li} f_{kl}(\mathbf{x}_i) \left(\mathbf{J}'_i - \Pi'_i \cdot \frac{\mathbf{U}_{kl}}{2}\right) \cdot \mathbf{r}_{kl}$$



Conceptually: useful to support DPD as a coarse-grained (CG) model

Practically: force fields are useless and can not reproduce MD system



Flekkoy&Coveney, PRL, 1999

Mori-Zwanzig Projection

Consider a canonical ensemble Γ . Def: A, B are two variables in Γ , noted by $A(\Gamma)$, $B(\Gamma)$. Def: Projection Operator P, Q

$$PB(\Gamma, t) = \frac{(B(\Gamma, t), A(\Gamma, t))}{(A(\Gamma, t), A(\Gamma, t))} A(\Gamma)$$
(1)

$$Q = 1 - P \tag{2}$$

Consider the time evolution operator e^{iLt} .

$$e^{iLt} = e^{iQLt} + \int_0^t d\tau e^{iQL(t-\tau)} iPLe^{iQL\tau}$$
(3)

The we have

$$\frac{dA(t)}{dt} = e^{iLt}iLA = e^{iLt}i(Q+P)LA \tag{4}$$

$$e^{iLt}iPLA = \frac{(iLA, A)}{(A, A)}e^{iLt}A = i\Omega A(t)$$
(5)

$$\frac{dA(t)}{dt} = i\Omega A(t) + e^{iLt} iQLA$$
$$= i\Omega A(t) + \int_0^t d\tau e^{iQL(t-\tau)} iPLe^{iQL\tau} iQLA + e^{iQLT} iQLA$$



Mori-Zwanzig Projection

Given A the coarse-grained velocity term, we identify $e^{iQLT}iQLA$ as the random force $\delta F(t).$ Since

$$(\delta F(t), A) = (e^{iQLt} iQLA, A) = (Q\delta F(t), A) = 0$$
(7)

$$iPLe^{iQLt}iQLA = iPL\delta F(t) = iPLQ\delta F(t)$$

$$= \frac{(iLQ\delta F(t), A)}{(A, A)}A = -\frac{(\delta F(t), iQLA)}{(A, A)}A$$

$$= -\frac{(\delta F(t), \delta F(0))}{(A, A)}A = -K(t)A$$
(8)

$$\begin{aligned} \frac{dA(t)}{dt} &= i\Omega A(t) - \int_0^t d\tau e^{iQL(t-\tau)} K(\tau) A + \delta F(t) \\ &= i\Omega A(t) - \int_0^t d\tau K(\tau) A(t-\tau) + \delta F(t) \end{aligned}$$



(9)

Mori-Zwanzig Projection

Specifically, if A(t) is the coarse-grained V(t), then

$$\frac{dV(t)}{dt} = i\Omega V(t) - \int_{0}^{t} d\tau \frac{\langle \delta F(\tau) \delta F(0) \rangle}{V(t)^{2}} V(t-\tau) + \delta F(t) \quad (10)$$

$$i\Omega V(t) = \frac{1}{\beta} \frac{\partial \ln \omega(\mathbf{R})}{\partial R} \quad (11)$$

$$\omega(\mathbf{R}) = \frac{\int d^{N} \hat{\mathbf{r}} \,\delta\left(\hat{\mathbf{R}} - \mathbf{R}\right) e^{-\beta U}}{\int d^{N} \hat{\mathbf{r}} \,e^{-\beta U}}, \quad (12)$$

Mori, ProgTheorPhys., 1965 Zwanzig, Oxford Uni. Press, 2001 Kinjo&Hyodo, PRE, 2007



MZ formalism as practical tool

Consider an atomistic system consisting of N atoms which are grouped into K clusters, and N_c atoms in each cluster. The Hamiltonian of the system is:

$$H = \sum_{\mu=1}^{K} \sum_{i=1}^{N_{C}} \frac{\mathbf{p}_{\mu,i}^{2}}{2m_{\mu,i}} + \frac{1}{2} \sum_{\mu,\nu} \sum_{i,j\neq i} V_{\mu i,\nu j}$$

Theoretically, the dynamics of the atomistic system can be mapped to a coarse-grained or mesoscopic level by using Mori-Zwanzig projection operators.

The equation of motion for coarse-grained particles can be written as: (in the following page)



MZ formalism as practical tool

Equation of motion for coarse-grained particles

1. Pairwise approximation: $\mathbf{F}_{\mu} \approx \sum_{\mu \neq \nu} \mathbf{F}_{\mu\nu}$ 2. Markovian approximation: $\langle \delta \mathbf{F}_{\mu}^{\vartheta}(t) \cdot \delta \mathbf{F}_{\nu}^{\vartheta}(0) \rangle = \Gamma_{\mu\nu} \cdot \delta(t)$





Lei, Caswell, &Karniadakis, PRE, 2010

Dynamical properties of constrained fluids

Mean square displacement (long time scale)



Coarse-graining unconstrained polymer melts

Natural bonds

 $N_c = 11$ $N_c = 21$ $N_c = 31$ $N_c = 11$ $N_c = 10$ $N_c = 10$ 1.0



 $\begin{aligned} & \textbf{WCA Potential} + \textbf{FENE Potential} \\ & V_{WCA}(r) = \begin{cases} 4\varepsilon \left[\left(\frac{\sigma}{r}\right)^{12} - \left(\frac{\sigma}{r}\right)^6 + \frac{1}{4} \right]; \ r \leq 2^{1/6}\sigma \\ 0; \ r > 2^{1/6}\sigma \end{cases} \\ & V_B(r) = \begin{cases} -\frac{1}{2}kR_0^2 \ln \left[1 - (r/R_0)^2 \right]; \ r \leq R_0 \\ \infty; \ r > R_0 \end{cases} \end{aligned}$

NVT ensemble with Nose-Hoover thermostat.



Directions for pairwise interactions between neighboring clusters



- 1. Parallel direction: $\mathbf{r}_{ij} = \mathbf{r}_i - \mathbf{r}_j$ $\mathbf{e}_{ij} = \mathbf{r}_{ij} / |r_{ij}|$
- 2. Perpendicular direction #1: $\mathbf{v}_{ij} = \mathbf{v}_i - \mathbf{v}_j$ $\mathbf{v}_{ij}^{\perp} = \mathbf{v}_{ij} - (\mathbf{v}_{ij} \cdot \mathbf{e}_{ij}) \cdot \mathbf{e}_{ij}$ $\mathbf{e}_{ij}^{\perp 1} = \mathbf{v}_{ij}^{\perp} / |v_{ij}^{\perp}|$ 3. Perpendicular direction #2:
 - $\mathbf{e}_{ij}^{\perp 2} = \mathbf{e}_{ij} \times \mathbf{e}_{ij}^{\perp 1}$



DPD force fields from MD simulation

Conservative

Dissipative (parallel one)



Li, Bian, Caswell, &Karniadakis, 2014



Performance of the MZ-DPD model ($N_c = 11$)

Quantities	MD	MZ-DPD (error)
Pressure	0.191	0.193 (+1.0%)
Diffusivity (Integral of VACF)	0.119	0.138 (+16.0%)
Viscosity	0.965	0.851 (-11.8%)
Schmidt number	8.109	6.167 (-23.9%)
Stokes-Einstein radius	1.155	1.129 (-2.2%)



Performance of the MZ-FDPD model ($N_c = 11$)

Quantities	MD	MZ-FDPD (error)
Pressure	0.191	0.193 (+1.0%)
Diffusivity (Integral of VACF)	0.119	0.120 (+ 0.8%)
Viscosity	0.965	0.954 (-1.1%)
Schmidt number	8.109	7.950 (-2.0%)
Stokes-Einstein radius	1.155	1.158 (+0.3%)



Conclusion&Outlook

- Invented by physics intuition
- Statistical physics on solid ground

 Flucutation-dissipation theorem
 Canonical ensemble (NVT)
- DDD < Novien Staker eque
- DPD <----> Navier-Stokes equations
- Coarse-graining microscopic system
 Mori-Zwanzig formalism

